

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, 15/62, 5/10, 1/21, C07K 14/515, 14/52	A1	(11) International Publication Number: WO 00/37642 (43) International Publication Date: 29 June 2000 (29.06.00)
(21) International Application Number: PCT/US99/30900 (22) International Filing Date: 23 December 1999 (23.12.99) (30) Priority Data: 60/113,387 23 December 1998 (23.12.98) US (71) Applicant (for all designated States except US): REGENERON PHARMACEUTICALS, INC. [US/US]; 777 Old Saw Mill River Road, Tarrytown, NY 10591-6707 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DAVIS, Samuel, J. [US/US]; 332 W. 88th Street, #B2, New York, NY 10024 (US). GALE, Nicholas, W. [US/US]; Apartment 46V, 177 White Plains Road, Tarrytown, NY 10591 (US). YANCOPOULOS, George, D. [US/US]; 1519 Baptist Church Road, Yorktown Heights, NY 10598 (US). STAHL, Neil [US/US]; RD # 10, Kent Shore Drive, Carmel, NY 10512 (US). (74) Agents: PALLADINO, Linda, O.; Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, NY 10591 (US) et al.		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHOD OF ENHANCING THE BIOLOGICAL ACTIVITY OF LIGANDS		
(57) Abstract Novel fusion polypeptide ligands that bind Eph family receptors or the Tie-2 receptor are identified, and methods for making the fusion polypeptide ligands in biologically active form are described. Nucleic acids encoding these novel fusion polypeptide ligands enable production of the fusion polypeptide ligands. The method of making the nucleic acids and the fusion polypeptide ligands is broadly applicable to the production of polypeptide ligands in general, resulting in improved affinity and/or increased activity of the ligand when compared to its native form.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHOD OF ENHANCING THE BIOLOGICAL ACTIVITY OF LIGANDS

This application claims priority of U.S. Application No. 60/113,387, filed December 23, 1998. Throughout this application, various publications are
5 cited. The disclosures of each and all of those publications are hereby incorporated by reference in their entireties into this application.

INTRODUCTION

10 The present invention provides for novel methods for producing novel fusion polypeptide ligands that have enhanced biological activity as compared to the polypeptide ligands in their native form. The invention also provides for nucleic acids useful for producing biologically active fusion polypeptide ligands, and the fusion polypeptide ligands themselves.

15

BACKGROUND OF THE INVENTION

The ability of polypeptide ligands to bind cells and thereby elicit a phenotypic response such as cell growth, survival or differentiation is often
20 mediated through transmembrane tyrosine kinases. The extracellular portion of each receptor tyrosine kinase (RTK) is generally the most distinctive portion of the molecule, as it provides the protein with its ligand-recognizing characteristic. Binding of a ligand to the extracellular domain results in signal transduction via an intracellular tyrosine kinase
25 catalytic domain which transmits a biological signal to intracellular target proteins. The particular array of sequence motifs of this cytoplasmic, catalytic domain determines its access to potential kinase substrates (Mohammadi, et al., 1990, Mol. Cell. Biol., 11: 5068-5078; Fantl, et al., 1992, Cell, 69:413-413).

30

RTKs appear to undergo dimerization or some related conformational change following ligand binding (Schlessinger, J., 1988, Trend Biochem. Sci.

- 13:443-447; Ullrich and Schlessinger, 1990, Cell, 61:203-212; Schlessinger and Ullrich, 1992, Neuron 9:383-391); molecular interactions between dimerizing cytoplasmic domains lead to activation of kinase function. In some instances, such as the growth factor platelet derived growth factor (PDGF), the ligand is a dimer that binds two receptor molecules (Hart, et al. , 5 1988, Science, 240: 1529-1531; Heldin, 1989, J. Biol. Chem. 264:8905-8912) while, for example, in the case of EGF, the ligand is a monomer (Weber, et al., 1984, J. Biol. Chem., 259:14631-14636).
- 10 Throughout the history of the biotechnology industry, many novel genes and associated proteins have been identified by virtue of their sequence homology with known genes. Many such proteins are purported to be receptors, but since their cognate ligands have not been identified, they are referred to as orphan receptors. The screening of many of these orphan 15 receptors often leads to the identification of ligands that are capable of binding to the receptor, although the binding is often not associated with activation of any intracellular kinases or any other phenotypic change. Such was the case for members of the Eph receptor family. For sake of clarity, applicants incorporate by reference herein a letter cited as Eph 20 Nomenclature Committee, 1997, published in Cell vol. 90: 403-403 (1997) which sets forth a nomenclature for the Eph Receptor and Eph Ligand Families.

- Little, if any, biological activity had been observed in response to binding of 25 a ligand to an Eph family receptor prior to the discovery as set forth in U.S. Patent No. 5,747,033 issued May 5, 1998. That patent describes the concept of "clustering" whereby the soluble domains of ligands were combined to create multimers capable of activating the cognate receptors. Applicants have now extended the concept of clustering to additional ligands outside 30 the Eph family, for example, the Tie-2 receptor ligands known as the angiopoietins, and have also discovered that this method for production of homogeneous forms of clustered ligands is broadly applicable to improve

the affinity and/or increase the activity of a ligand as compared to the native form of the ligand.

Angiopoietin-1 (Ang) is one of two known ligands for the Tie-2 receptor and has been shown to be an agonist for Tie-2 (Davis, et al, 1996, Cell 87:1161-1169), whereas the second known ligand, angiopoietin-2, has been shown to be a naturally occurring antagonist of the Tie-2 receptor (Maisonpierre, et al., 1997, Science 277:55-60). Ang1* is a mutant form of angiopoietin-1 that comprises the N-terminal domain of angiopoietin-2 fused to the coiled-coil domain and the fibrinogen domain of angiopoietin-1 and that has a Cys to Ser mutation at amino acid 245. Ang1* has been shown to be a potent agonist for the Tie-2 receptor.

Experiments with mutants of angiopoietin-1 and angiopoietin-2 have demonstrated that the fibrinogen domains (FD) are the receptor-binding domains, and that dimerized versions of, for example Ang-1-FD-Fc, (i.e., the fibrinogen domain of Ang-1 fused to an Fc domain), can bind to the Tie-2 receptor with much higher affinity than monomeric Ang-1-FD (dimerization occurs due to the interaction between the Fc components of adjacent molecules). However, Ang-1-FD-Fc is not able to induce phosphorylation (activate) the Tie-2 receptor on endothelial cells unless it is further clustered with goat anti-human Fc antibodies (Jackson Immunoresearch). For this reason, mutant versions of Ang-1-FD and Ang-2-FD (i.e., the fibrinogen domain of Ang-2) were designed that were intrinsically more highly clustered.

SUMMARY OF THE INVENTION

The present invention provides for novel, biologically active, soluble forms of polypeptide ligands that bind to receptors on cells. Such polypeptide ligands are useful in promoting a differential function and/or influencing the phenotype, such as growth and/or proliferation, of receptor-bearing

cells. The invention also provides for nucleic acids encoding such polypeptide ligands, and both prokaryotic and eukaryotic expression systems for producing such polypeptide ligands. According to the invention, soluble forms of the polypeptide ligands described herein may be used to promote biological responses in receptor-expressing cells. In particular, a general method is described herein which produces fusion polypeptide ligands that may then be clustered, which functions to make otherwise inactive soluble polypeptide ligands biologically active, or which enhances the biological activity of polypeptide ligands that, absent such clustering, would have lower levels of biological activity. This method may be used to cluster a plurality of (more than one) receptor binding domains from any ligand which has improved affinity and/or increased activity (i.e. signaling ability) when clustered as compared to the native form of the ligand.

DESCRIPTION OF THE FIGURES

Figure 1A-1E - Nucleic acid sequence and deduced amino acid sequence of Ang-1-FD-FD-Fc.

Figure 2A-2E - Nucleic acid sequence and deduced amino acid sequence of Ang-2-FD-FD-Fc.

Figure 3A-3E - Nucleic acid sequence and deduced amino acid sequence of Ang-1-FD-Fc-FD.

Figure 4A-4E - Nucleic acid sequence and deduced amino acid sequence of Ang-2-FD-Fc-FD.

Figure 5 - Molecular Weight Analysis of Ang-1-FD-Fc-FD protein. SDS PAGE analyses showing a band running at about 210kD under non-reducing conditions (lane 3) and a band running at about 85kD under reducing conditions (lane 7).

Figure 6 - Light scatter analysis to confirm the molecular weight of Ang-1-FD-Fc-FD and to determine whether or not the protein is a homogeneous species. Light scattering is a function of mass and concentration of a macromolecule. To determine molecular weight, the protein sample was injected onto a gel filtration column and the effluent is monitored with an on line light scattering detector and a refractive index and/or a UV detector. The on line refractive index detector or UV detector serve to measure protein concentration. Astra 4.7 Software (Wyatt Technology Corporation, Santa Barbara, CA) is used to calculate the protein concentration. The molecular weight of protein is then calculated from the angular dependence of light scattering. The molecular weight of the dimeric protein appears to be approximately 200kD and presence of a single peak implies that the protein solution is homogenous.

Figure 7 - Molecular Weight Analysis of Ang-2-FD-Fc-FD. SDS PAGE analyses showing a band running at about 200kD under non-reducing conditions (lanes 7 and 8) and a band running at about 88kD under reducing conditions (lanes 3 and 4).

Figure 8 - Light scatter analysis to confirm the molecular weight of Ang-2-FD-Fc-FD and to determine whether or not the protein is a homogeneous species. Light scattering is a function of mass and concentration of a macromolecule. To determine molecular weight, the protein sample was injected onto a gel filtration column and the effluent is monitored with an on line light scattering detector and a refractive index and/or a UV detector. The on line refractive index detector or UV detector serve to measure protein concentration. Astra 4.7 Software (Wyatt Technology Corporation, Santa Barbara, CA) is used to calculate the protein concentration. The molecular weight of protein is then calculated from the angular dependence of light scattering. The molecular weight of the dimeric protein appears to be approximately 171kD and presence of a single peak implies that the

protein solution is homogenous.

Figure 9 - Ang1*-mediated vs. Ang-1-FD-Fc-FD-mediated Tie-2 receptor phosphorylation in EAhy926 cells. A standard phosphorylation assay
5 revealed that Ang-1-FD-Fc-FD was equivalent to Ang1* in its ability to stimulate phosphorylation of the Tie-2 receptor in EAhy926.

Figure 10 - Ability of Ang-2-FD-Fc-FD to block Ang1*-mediated Tie-2 receptor phosphorylation in EAhy926 cells. In a standard phosphorylation
10 assay, Ang-2-FD-Fc-FD is able to block Ang1* stimulation of the Tie-2 receptor when it is present in at least a 10-15 fold molar excess of Ang1*.

Figure 11 - Ability of angiopoietin-2 to block Ang1*-mediated Tie-2 receptor phosphorylation in EAhy926 cells. In a standard phosphorylation assay, at a
15 20 fold molar excess, angiopoietin-2 is not able to reduce the Ang1*-mediated phosphorylation level to 50%. This result, coupled with the results described in Figure 10 implies that Ang-2-FD-Fc-FD is a more potent inhibitor of Ang1*-mediated Tie-2 receptor phosphorylation than angiopoietin-2.

20

Figure 12 - Ability of Ang-2-FD-Fc-FD to block angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in EAhy926 cells. In a standard phosphorylation assay, it is shown that while there is a trend toward blocking angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in
25 these cells, Ang-2-FD-Fc-FD seems to be more effective at blocking Ang1*-mediated phosphorylation of Tie-2, as shown in Figure 10.

Figure 13 - Ability of angiopoietin-2 to block angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in EAhy926 cells. In a standard
30 phosphorylation assay, it is shown that there is a trend toward blocking angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in these cells, but, like Ang-2-FD-Fc-FD, angiopoietin-2 seems to be more effective at

blocking Ang1*-mediated phosphorylation of Tie-2, as shown in Figure 11.

Figure 14A-14E - Nucleic acid sequence and deduced amino acid sequence of Ephrin-B1-Ephrin-B1-Fc.

5

Figure 15A-15E - Nucleic acid sequence and deduced amino acid sequence of Ephrin-B2-Ephrin-B2-Fc.

Figure 16 - Comparison of Ephrin-B1-Fc, Ephrin-B1-Ephrin-B1-Fc, Ephrin-B2-Fc and Ephrin-B2-Ephrin-B2-Fc in standard EphB2 phosphorylation assays. COS cells were serum-starved and then left untreated (UT), lane 1, or were treated with unclustered and clustered Ephrin-B1-Fc (Efn-B1), lanes 2 and 3. COS cells were also treated with unclustered and clustered Ephrin-B1-Ephrin-B1-Fc (Efn-B1 DD), lanes 4 and 5. In addition cells were likewise
10 treated with unclustered and clustered Ephrin-B2-Fc (Efn-B2), lanes 6 and 7 and with unclustered and clustered Ephrin-B2-Ephrin-B2-Fc (Efn-B2 DD), lanes 8 and 9. The extent of EphB2 phosphorylation was assessed by anti-phosphotyrosine western blotting (upper panels) and the relative amounts of EphB2 in each lane was determined by anti-EphB2 western blotting
15 (lower panels).
20

Figure 17 - Ang1*-mediated vs. stable CHO clone-derived Ang-1-FD-Fc-FD-mediated Tie-2 receptor phosphorylation in EAhy926 cells. EAhy926 cells were stimulated with 0.4 µg/ml Ang1* or 0.2 µg/ml or 0.4 µg/ml stable
25 CHO clone-derived Ang-1-FD-Fc-FD protein. A standard phosphorylation assay revealed that stable CHO clone-derived Ang-1-FD-Fc-FD was equivalent to Ang1* in its ability to stimulate phosphorylation of the Tie-2 receptor in EAhy926 cells.

30 Figure 18 - Ability of stable CHO clone-derived Ang-2-FD-Fc-FD to block stable CHO clone-derived Ang-1-FD-Fc-FD-mediated Tie-2 receptor phosphorylation in EAhy926 cells. EAhy926 cells were treated with 0.2

µg/ml of the Tie-2 agonist Ang-1-FD-Fc-FD and 2 µg/ml, 4 µg/ml, 8 µg/ml or 16 µg/ml of stable CHO clone-derived Ang-2-FD-Fc-FD. Ang-2-FD-Fc-FD is able to block or stable CHO clone-derived Ang-1-FD-Fc-FD stimulation of the Tie-2 receptor when it is present in at least a 40 fold molar excess of
5 stable CHO clone-derived Ang-1-FD-Fc-FD.

DETAILED DESCRIPTION OF THE INVENTION

As described in greater detail below, applicants have discovered a method
10 for "clustering" polypeptide ligands, which functions to make otherwise inactive soluble polypeptide ligands biologically active, or which enhances the biological activity of polypeptide ligands that, absent such clustering, would have lower levels of biological activity. This method may be used to cluster a plurality of (more than one) receptor binding domains from any
15 ligand which has improved affinity and/or increased activity (i.e. signaling ability) when clustered as compared to the native form of the ligand.

The present invention provides for a nucleic acid encoding a fusion polypeptide wherein the fusion polypeptide comprises a first subunit
20 comprising at least one copy of the receptor binding domain of a ligand, the first subunit being fused to the N-terminal end of a multimerizing component, said multimerizing component being fused at its C-terminal end to a second subunit comprising at least one copy of the receptor binding domain of a ligand.

25 In one embodiment of the invention, the receptor binding domains of the first and second subunits are copies of the receptor binding domain of the same ligand. The first and second subunits may each have one or more than one copy of the receptor binding domain of the ligand. In specific
30 embodiments of the invention, the receptor binding domain is the fibrinogen domain of angiopoietin-1 or angiopoietin-2. Alternatively, the

receptor binding domain is from a ligand selected from the group consisting of the EPH family of ligands (i.e., the ephrins).

In another embodiment of the invention, the receptor binding domains of the first subunit are copies of the receptor binding domain of a different ligand from the receptor binding domains of the second subunit. For example, the first subunit may comprise the receptor binding domain of an angiopoietin and the second subunit may comprise the receptor binding domain of vascular endothelial growth factor (VEGF). Alternatively, the first subunit may comprise the receptor binding domain of VEGF and the second subunit may comprise the receptor binding domain an angiopoietin. Still further, the first and second subunits may each have one or more than one copy of the receptor binding domain of their respective ligand.

By "receptor binding domain" what is meant is the minimal portion of the ligand that is necessary to bind its receptor.

In preferred embodiments of the invention, the multimerizing component comprises an immunoglobulin derived domain. More specifically, the immunoglobulin derived domain may be selected from the group consisting of the Fc domain of IgG, the heavy chain of IgG, and the light chain of IgG. In another embodiment, the multimerizing component may be an Fc domain from which the first five amino acids (including a cysteine) have been removed to produce a multimerizing component referred to as Fc(Δ C1).

The present invention also provides for fusion polypeptides encoded by the nucleic acid molecules of the invention. Preferably, the fusion polypeptides are in multimeric form, due to the function of the multimerizing component. In a preferred embodiment, the multimer is a dimer. Suitable multimerizing components are described in European Patent Application of ZymoGenetics, Inc., Publication No. EP 0 721 983 A1 published 17 July 1996

and include S. cerevisiae repressible acid phosphatase (Mizunaga et al., 1988, J. Biochem. (Tokyo) 103:321-326); the S. cerevisiae type 1 killer preprotoxin (Sturley et al., 1986, EMBO J. 5:3381-3390); the S. calbergensis alpha galactosidase melibiase (Sumner-Smith, et al., 1985, Gene 36:333-340); and
5 the Neurospora crassa ornithine decarboxylase (Digangi, et al., 1987, J. Biol. Chem. 262:7889-7893). Sequences encoding an immunoglobulin heavy chain hinge region (Takahashi et al., 1982, Cell 29:671-679); the S. cerevisiae SUC2 gene (Carlson et al., 1983, Mol. Cell. Biol. 3:439-447); immunoglobulin
10 gene sequences, and portions thereof. In a preferred embodiment of the invention, immunoglobulin gene sequences, especially one encoding the Fc domain, are used to encode the multimerizing component.

The present invention also contemplates a vector which comprises the nucleic acid molecule of the invention as described herein.

15 Also provided is an expression vector comprising a nucleic acid molecule of the invention as described herein, wherein the nucleic acid molecule is operatively linked to an expression control sequence. Also provided is a host-vector system for the production of a fusion polypeptide which
20 comprises the expression vector of the invention which has been introduced into a host cell suitable for expression of the fusion polypeptide. The suitable host cell may be a bacterial cell such as E. coli, a yeast cell, such as Pichia pastoris, an insect cell, such as Spodoptera frugiperda, or a mammalian cell, such as a COS or CHO cell.

25 The present invention also provides for methods of producing the fusion polypeptides of the invention by growing cells of the host-vector systems described herein, under conditions permitting production of the fusion polypeptide and recovering the fusion polypeptide so produced.

30 The fusion polypeptides useful for practicing the present invention may be prepared by expression in a prokaryotic or eukaryotic expression system.

The recombinant gene may be expressed and the polypeptide purified utilizing any number of methods. The gene may be subcloned into a bacterial expression vector, such as for example, but not by way of limitation, pCP110.

5

The fusion polypeptides may be purified by any technique which allows for the subsequent formation of a stable, biologically active protein. For example, and not by way of limitation, the factors may be recovered from cells either as soluble proteins or as inclusion bodies, from which they may be extracted quantitatively by 8M guanidinium hydrochloride and dialysis. In order to further purify the factors, conventional ion exchange chromatography, hydrophobic interaction chromatography, reverse phase chromatography or gel filtration may be used.

15 The present invention also provides for a nucleic acid encoding a fusion polypeptide wherein the fusion polypeptide comprises more than one copy of the receptor binding domain of a ligand in tandem, and wherein either the N-terminal or the C-terminal receptor binding domain is also fused to a multimerizing component. In one embodiment of the invention, the
20 receptor binding domains are fused contiguously. In another embodiment of the invention, the receptor binding domains are from a ligand that is not a member of the EPH family of ligands (i.e., not an ephrin). In specific embodiments of the invention, the receptor binding domain is the fibrinogen domain of angiopoietin-1 or angiopoietin-2. In an alternative
25 embodiment, the receptor binding domain is from vascular endothelial growth factor (VEGF). In another embodiment, the receptor binding domain is from an ephrin.

By "receptor binding domain" what is meant is the minimal portion of the
30 ligand that is necessary to bind its receptor.

In preferred embodiments of the invention, the multimerizing component comprises an immunoglobulin derived domain. More specifically, the immunoglobulin derived domain may be selected from the group consisting of the Fc domain of IgG, the heavy chain of IgG, and the light chain of IgG. In another embodiment, the multimerizing component may be an Fc domain from which the first five amino acids (including a cysteine) have been removed to produce a multimerizing component referred to as Fc(Δ C1).

The present invention also provides for fusion polypeptides encoded by the nucleic acid molecules of the invention. Preferably, the fusion polypeptides are in multimeric form, due to the function of the multimerizing component. In a preferred embodiment, the multimer is a dimer. Suitable multimerizing components are described in European Patent Application of ZymoGenetics, Inc., Publication No. EP 0 721 983 A1 published 17 July 1996 and include *S. cerevisiae* repressible acid phosphatase (Mizunaga et al., 1988, J. Biochem. (Tokyo) 103:321-326); the *S. cerevisiae* type 1 killer preprotoxin (Sturley et al., 1986, EMBO J. 5:3381-3390); the *S. calbergensis* alpha galactosidase melibiase (Sumner-Smith, et al., 1985, Gene 36:333-340); and the *Neurospora crassa* ornithine decarboxylase (Digangi, et al., 1987, J. Biol. Chem. 262:7889-7893). Sequences encoding an immunoglobulin heavy chain hinge region (Takahashi et al., 1982, Cell 29:671-679); the *S. cerevisiae* SUC2 gene (Carlson et al., 1983, Mol. Cell. Biol. 3:439-447); immunoglobulin gene sequences, and portions thereof. In a preferred embodiment of the invention, immunoglobulin gene sequences, especially one encoding the Fc domain, are used to encode the multimerizing component.

The present invention also contemplates a vector which comprises the nucleic acid molecule of the invention as described herein.

Also provided is an expression vector comprising a nucleic acid molecule of the invention as described herein, wherein the nucleic acid molecule is

operatively linked to an expression control sequence. Also provided is a host-vector system for the production of a fusion polypeptide which comprises the expression vector of the invention which has been introduced into a host cell suitable for expression of the fusion polypeptide.

- 5 The suitable host cell may be a bacterial cell such as E. coli, a yeast cell, such as Pichia pastoris, an insect cell, such as Spodoptera frugiperda, or a mammalian cell, such as a COS or CHO cell.

- 10 The present invention also provides for methods of producing the fusion polypeptides of the invention by growing cells of the host-vector systems described herein, under conditions permitting production of the fusion polypeptide and recovering the fusion polypeptide so produced.

- 15 The fusion polypeptides useful for practicing the present invention may be prepared by expression in a prokaryotic or eukaryotic expression system. The recombinant gene may be expressed and the polypeptide purified utilizing any number of methods. The gene may be subcloned into a bacterial expression vector, such as for example, but not by way of limitation, pCP110.

- 20 The fusion polypeptides may be purified by any technique which allows for the subsequent formation of a stable, biologically active protein. For example, and not by way of limitation, the factors may be recovered from cells either as soluble proteins or as inclusion bodies, from which they may be extracted quantitatively by 8M guanidinium hydrochloride and dialysis. In order to further purify the factors, conventional ion exchange chromatography, hydrophobic interaction chromatography, reverse phase chromatography or gel filtration may be used.

- 30 The Examples describe the preparation of novel polypeptide ligands that comprise a receptor binding domain of a member of the Eph (Eph transmembrane tyrosine kinase family ligands) family of ligands or of a

member of the angiopoietin family of ligands that can bind the Tie-2 receptor.

- For a description of novel Eph family ligands, methods of making and using them, as well as the sequences of EHK-1L, B61 and ELK-L, together with a description of a method of enhancing the biological activity of EPH family ligands by clustering them, applicants refer to U.S. Patent No. 5,747,033 issued on May 5, 1998 which is hereby incorporated by reference in its entirety. Applicants further refer to International Application PCT/US93/10879, published as WO 94/11020 on May 26, 1994; and International Application PCT/US96/17201 published as WO 97/15667 entitled "Biologically Active EPH Family Ligands" each of which is hereby incorporated by reference in its entirety.
- As has been previously reported, a family of ligands for the TIE-2 receptor has been discovered and named the Angiopoietins. This family, consisting of TIE-2 ligand 1 (Ang-1); TIE-2 ligand 2 (Ang-2); TIE ligand 3 (Ang-3); and TIE ligand 4 (Ang-4) has been extensively characterized. For a description of the cloning, sequencing and characterization of the angiopoietins, as well as for methods of making and uses thereof, including the production and characterization of modified and chimeric ligands thereof, reference is hereby made to the following publications, each of which is incorporated by reference herein in its entirety: U.S. Patent No. 5,521,073 issued May 28, 1996; U.S. Patent No. 5,643,755 issued July 1, 1997; U.S. Patent No. 5,650,490 issued July 22, 1997; U.S. Patent No. 5,814,464 issued September 29, 1998; U.S. Patent No. 5,879,672 issued March 9, 1999; U.S. Patent No. 5,851,797 issued December 22, 1998; PCT International Application entitled "TIE-2 Ligands Methods of Making and Uses Thereof," published as WO 96/11269 on 18 April 1996 in the name of Regeneron Pharmaceuticals, Inc.; PCT International Application entitled "TIE-2 Ligands Methods of Making and Uses Thereof," published as WO 96/31598 on 10 October 1996 in the name of Regeneron Pharmaceuticals, Inc.; PCT International Application entitled

"TIE-2 Receptor Ligands (TIE Ligand-3; TIE Ligand-4) And Their Uses," published as WO 97/48804 on 24 December 1997 in the name of Regeneron Pharmaceuticals, Inc; and PCT International Application entitled "Modified TIE-2 Receptor Ligands," published as WO 98/05779 on 12 February 1998 in the name of Regeneron Pharmaceuticals, Inc.

When used herein, fusion polypeptide includes functionally equivalent molecules in which amino acid residues are substituted for residues within the sequence resulting in a silent or conservative change. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent or conservative alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Also included within the scope of the invention are proteins or fragments or derivatives thereof which exhibit the same or similar biological activity and derivatives which are differentially modified during or after translation, e.g., by glycosylation, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc.

Cells that express the fusion polypeptides of the invention are genetically engineered to produce them by, for example, transfection, transduction, electroporation, or microinjection.

The present invention encompasses the nucleic acid sequences encoding the fusion polypeptides of the invention, as well as sequences that hybridize under stringent conditions to nucleic acid sequences that are
5 complementary to the nucleic acid sequences of the invention. Stringent conditions are set forth in, for example, Sambrook, et al. Molecular Cloning: A Laboratory Manual, 2 ed. Vol. 1, pp. 101-104, Cold Spring Harbor Laboratory Press (1989). In addition, the present invention encompasses nucleic acid sequences that are different from the nucleic acid sequences of
10 the invention but which nevertheless encode the fusion polypeptides of the invention due to the degeneracy of the genetic code.

In addition, the present invention contemplates use of the fusion polypeptides described herein in tagged forms.

15 Any of the methods known to one skilled in the art for the insertion of DNA fragments into a vector may be used to construct expression vectors encoding the fusion polypeptides of the invention using appropriate transcriptional/translational control signals and the protein coding
20 sequences. These methods may include in vitro recombinant DNA and synthetic techniques and in vivo recombinations (genetic recombination). Expression of nucleic acid sequence encoding the fusion polypeptides of the invention may be regulated by a second nucleic acid sequence so that the fusion polypeptide is expressed in a host transformed with the recombinant
25 DNA molecule. For example, expression of the fusion polypeptides described herein may be controlled by any promoter/enhancer element known in the art. Promoters which may be used to control expression of the fusion polypeptide include, but are not limited to the long terminal repeat as described in Squinto et al., (1991, Cell 65:1-20); the SV40 early
30 promoter region (Bernoist and Chambon, 1981, Nature 290:304-310), the CMV promoter, the M-MuLV 5' terminal repeat the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980,

Cell 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:144-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, Nature 296:39-42); prokaryotic expression vectors such as the b-lactamase promoter (Villa-Kamaroff, et al., 5 1978, Proc. Natl. Acad. Sci. U.S.A. 75:3727-3731), or the *tac* promoter (DeBoer, et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:21-25), see also "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242:74-94; promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADH (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) 10 promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, Cell 38:639-646; Ornitz et al., 1986, Cold Spring Harbor Symp. Quant. Biol. 50:399-409; MacDonald, 15 1987, Hepatology 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, Nature 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, Cell 38:647-658; Adames et al., 1985, Nature 318:533-538; Alexander et al., 1987, Mol. Cell. Biol. 7:1436-1444), mouse mammary tumor virus control 20 region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, Cell 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, Genes and Devel. 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, Mol. Cell. Biol. 5:1639-1648; Hammer et al., 1987, Science 235:53-58); alpha 1-antitrypsin 25 gene control region which is active in the liver (Kelsey et al, 1987, Genes and Devel. 1:161-171), beta-globin gene control region which is active in myeloid cells (Mogram et al., 1985, Nature 315:338-340; Kollias et al., 1986, Cell 46:89-94); myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, Cell 48:703-712); 30 myosin light chain-2 gene control region which is active in skeletal muscle (Shani, 1985, Nature 314:283-286), and gonadotropic releasing hormone gene

control region which is active in the hypothalamus (Mason et al., 1986, Science 234:1372-1378).

Thus, according to the invention, expression vectors capable of being
5 replicated in a bacterial or eukaryotic host comprising Eph fusion
polypeptide encoding or angiopoietin fusion polypeptide encoding nucleic
acids as described herein, are used to transfect the host and thereby direct
expression of such nucleic acid to produce fusion polypeptides which may
then be recovered in biologically active form. As used herein, a biologically
10 active form includes a form capable of binding to the relevant receptor and
causing a differentiated function and/or influencing the phenotype of the
cell expressing the receptor. Such biologically active forms would, for
example, induce phosphorylation of the tyrosine kinase domain of the Etk-
1, Elk, or Tie2 receptor, or stimulation of synthesis of cellular DNA.

15 Expression vectors containing the nucleic acid inserts can be identified by
three general approaches: (a) DNA-DNA hybridization, (b) presence or
absence of "marker" gene functions, and (c) expression of inserted
sequences. In the first approach, the presence of a foreign nucleic acids
20 inserted in an expression vector can be detected by DNA-DNA hybridization
using probes comprising sequences that are homologous to an inserted
nucleic acid sequences. In the second approach, the recombinant
vector/host system can be identified and selected based upon the presence
or absence of certain "marker" gene functions (e.g., thymidine kinase
25 activity, resistance to antibiotics, transformation phenotype, occlusion body
formation in baculovirus, etc.) caused by the insertion of foreign nucleic
acid sequences in the vector. For example, if an efl nucleic acid sequence is
inserted within the marker gene sequence of the vector, recombinants
containing the insert can be identified by the absence of the marker gene
30 function. In the third approach, recombinant expression vectors can be
identified by assaying the foreign nucleic acid product expressed by the
recombinant. Such assays can be based, for example, on the physical or

functional properties of the nucleic acid product of interest, for example, by binding of a ligand to a receptor or portion thereof which may be tagged with, for example, a detectable antibody or portion thereof or binding to antibodies produced against the protein of interest or a portion thereof.

5

Cells of the present invention may transiently or, preferably, constitutively and permanently express the ephrin or angiopoietin fusion polypeptide as described herein.

10 The ephrin fusion polypeptides of the invention may be useful in methods of treating a patient suffering from a neurological disorder comprising treating the patient with an effective amount of the ephrin fusion polypeptide.

15 For example, the Elk receptor is expressed primarily in brain. Accordingly, it is believed that an Elk binding ephrin fusion polypeptide ligand will support the induction of a differential function and/or influence the phenotype, such as growth and/or survival of neural cells that express this receptor.

20

The present invention also provides for pharmaceutical compositions comprising the ephrin fusion polypeptide in a suitable pharmacologic carrier. The compositions may be administered systemically or locally. Any appropriate mode of administration known in the art may be used,
25 including, but not limited to, intravenous, intrathecal, intraarterial, intranasal, oral, subcutaneous, intraperitoneal, or by local injection or surgical implant. Sustained release formulations are also provided for.

As our understanding of neurodegenerative disease/neurotrauma becomes
30 clearer, it may become apparent that it would be beneficial to decrease the effect of endogenous Efl-6. Therefore, in areas of nervous system trauma, it may be desirable to provide Efl-6 antagonists, including, but not limited to,

fusion polypeptide forms of Efl-6 which may compete with cell-bound ligand for interaction with Elk receptor. It may be desirable to provide such antagonists locally at the injury site rather than systemically. Use of an Efl-6 antagonist providing implant may be desirable.

5

Alternatively, certain conditions may benefit from an increase in Efl-6 responsiveness. It may therefore be beneficial to increase the number or binding affinity of Efl-6 in patients suffering from such conditions.

10 The invention herein further provides for the development of a fusion polypeptide, as a therapeutic for the treatment of patients suffering from disorders involving cells, tissues or organs which express the TIE-2 receptor. Such molecules may be used in a method of treatment of the human or animal body, or in a method of diagnosis.

15

Because TIE-2 receptor has been identified in association with endothelial cells and, as was previously demonstrated, blocking of agonists of the receptor such as TIE-2 ligand 1 (Ang-1) has been shown to prevent vascularization, applicants expect that TIE-2 agonist fusion polypeptides of
20 the invention may be useful for the induction of vascularization in diseases or disorders where such vascularization is indicated. Such diseases or disorders would include wound healing, ischemia and diabetes. The ligands may be tested in animal models and used therapeutically as described for other agents, such as vascular endothelial growth factor
25 (VEGF), another endothelial cell-specific factor that is angiogenic. Ferrara, et al. U.S. Patent No. 5,332,671 issued July 26, 1994. The Ferrara reference, as well as other studies, describe in vitro and in vivo studies that may be used to demonstrate the effect of an angiogenic factor in enhancing blood flow to ischemic myocardium, enhancing wound healing, and in other therapeutic
30 settings wherein neoangiogenesis is desired. [see Sudo, et al., European Patent Application 0 550 296 A2 published July 7, 1993; Banai, et al. Circulation 89:2183-2189 (1994); Unger, et al. Am. J. Physiol. 266:H1588-H1595

(1994); Lazarous, et al. Circulation 91:145-153 (1995)]. According to the invention, the agonist fusion polypeptides may be used alone or in combination with one or more additional pharmaceutically active compounds such as, for example, VEGF or basic fibroblast growth factor (bFGF).

Conversely, antagonists of the TIE-2 receptor, such as TIE-2 receptorbodies or TIE-2 ligand 2 (Ang-2) as described in Example 9 in International Publication No. WO 96/31598 published 10 October 1996, have been shown to prevent or attenuate vascularization, and are thus expected to be useful in preventing or attenuating, for example, tumor growth. Similarly then, TIE-2 antagonist fusion polypeptides of the invention would also be useful for those purposes. These antagonists may be used alone or in combination with other compositions, such as anti-VEGF antibodies, that have been shown to be useful in treating conditions in which the therapeutic intent is to block angiogenesis.

For example, applicants have determined that TIE-2 ligands are expressed in cells within, or closely associated with, tumors. For example, TIE-2 ligand 2 (Ang-2) appears to be tightly associated with tumor endothelial cells. Accordingly, TIE-2 antagonist fusion polypeptides of the invention may also be useful in preventing or attenuating, for example, tumor growth.

In other embodiments, the TIE-2 agonist fusion polypeptides of the invention described herein may be used as hematopoietic factors. A variety of hematopoietic factors and their receptors are involved in the proliferation and/or differentiation and/or migration of the various cells types contained within blood. Because the TIE-2 receptors are expressed in early hematopoietic cells, the TIE-2 ligands are expected to play a comparable role in the proliferation or differentiation or migration of these cells. Thus, for example, TIE-2 agonist fusion polypeptide compositions may be prepared, assayed, examined in *in vitro* and *in vivo* biological systems and

used therapeutically as described in any of the following: Sousa, U.S. Patent No. 4,810,643, Lee, et al., Proc. Natl. Acad. Sci. USA 82:4360-4364 (1985) Wong, et al. Science, 228:810-814 (1985); Yokota, et al. Proc. Natl. Acad. Sci (USA) 81:1070 (1984); Bosselman, et al. WO 9105795 published May 2, 1991
5 entitled "Stem Cell Factor" and Kirkness, et al. WO 95/19985 published July 27, 1995 entitled "Haemopoietic Maturation Factor". Accordingly, the fusion polypeptides may be used to diagnose or treat conditions in which normal hematopoiesis is suppressed, including, but not limited to anemia, thrombocytopenia, leukopenia and granulocytopenia. In a preferred
10 embodiment, the fusion polypeptides may be used to stimulate differentiation of blood cell precursors in situations where a patient has a disease, such as acquired immune deficiency syndrome (AIDS) which has caused a reduction in normal blood cell levels, or in clinical settings in which enhancement of hematopoietic populations is desired, such as in
15 conjunction with bone marrow transplant, or in the treatment of aplasia or myelosuppression caused by radiation, chemical treatment or chemotherapy.

The fusion polypeptides of the present invention may be used alone, or in
20 combination with another pharmaceutically active agents such as, for example, cytokines, neurotrophins, interleukins, etc. In a preferred embodiment, the fusion polypeptides may be used in conjunction with any of a number of factors which are known to induce stem cell or other hematopoietic precursor proliferation, or factors acting on later cells in the
25 hematopoietic pathway, including, but not limited to, hemopoietic maturation factor, thrombopoietin, stem cell factor, erythropoietin, G-CSF, GM-CSF, etc.

In an alternative embodiment, TIE-2 receptor antagonist fusion
30 polypeptides are used to diagnose or treat patients in which the desired result is inhibition of a hematopoietic pathway, such as for the treatment of myeloproliferative or other proliferative disorders of blood forming organs

such as thrombocythemias, polycythemias and leukemias. In such embodiments, treatment may comprise use of a therapeutically effective amount of the fusion polypeptides as described herein.

- 5 Effective doses useful for treating these or other diseases or disorders may be determined using methods known to one skilled in the art [see, for example, Fingl, et al., *The Pharmacological Basis of Therapeutics*, Goodman and Gilman, eds. Macmillan Publishing Co., New York, pp. 1-46 ((1975))]. Pharmaceutical compositions for use according to the invention include the
- 10 fusion polypeptides described above in a pharmacologically acceptable liquid, solid or semi-solid carrier, linked to a carrier or targeting molecule (e.g., antibody, hormone, growth factor, etc.) and/or incorporated into liposomes, microcapsules, and controlled release preparation prior to administration *in vivo*. For example, the pharmaceutical composition may
- 15 comprise a fusion polypeptide in an aqueous solution, such as sterile water, saline, phosphate buffer or dextrose solution. Alternatively, the active agents may be comprised in a solid (e.g. wax) or semi-solid (e.g. gelatinous) formulation that may be implanted into a patient in need of such treatment. The administration route may be any mode of administration
- 20 known in the art, including but not limited to intravenously, intrathecally, subcutaneously, by injection into involved tissue, intraarterially, intranasally, orally, or via an implanted device.

- Administration may result in the distribution of the active agent of the
- 25 invention throughout the body or in a localized area. For example, in some conditions which involve distant regions of the nervous system, intravenous or intrathecal administration of agent may be desirable. In some situations, an implant containing active agent may be placed in or near the lesioned area. Suitable implants include, but are not limited to,
- 30 gelfoam, wax, or microparticle-based implants.

The present invention also provides for pharmaceutical compositions comprising the fusion polypeptides described herein, in a pharmacologically acceptable vehicle. The compositions may be administered systemically or locally. Any appropriate mode of administration known in the art may be used, including, but not limited to, intravenous, intrathecal, intraarterial, intranasal, oral, subcutaneous, intraperitoneal, or by local injection or surgical implant. Sustained release formulations are also provided for.

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

EXAMPLES

Angiopoietin ligands:

As described *supra*, experiments with mutants of Ang-1 and Ang-2 have demonstrated that the fibrinogen domains (FD) are the receptor-binding domains, and that dimerized versions (dimerization occurs due to the interaction between the Fc components of adjacent molecules), for example Ang-1-FD-Fc, can bind to the Tie-2 receptor with much higher affinity than monomeric Ang-1-FD. However, Ang-1-FD-Fc is not able to induce phosphorylation (activate) the Tie-2 receptor on endothelial cells unless it is further clustered with goat anti-human Fc antibodies (Jackson Immunoresearch). For this reason, mutant versions of Ang-1-FD and Ang-2-FD were designed that were intrinsically more highly clustered.

Two general types of nucleic acid molecules were constructed. The first type consisted of two tandem copies of Ang-1-FD fused to an Fc tag, thus leading

to a secreted polypeptide molecule that is dimeric with respect to the Fc tag but tetrameric with respect to Ang-1-FD. Similarly, two tandem copies of Ang-2-FD fused to an Fc tag constituted the angiopoietin-2 version of this type of construct. These molecules were designated Ang-1-FD-FD-Fc and
5 Ang-2-FD-FD-Fc, respectively.

In the second type of nucleic acid molecule constructed, two copies of Ang-1-FD were connected by an Fc tag bridging between them, thus creating the structure Ang-1-FD-Fc-FD that is still dimeric with respect to the Fc, as well
10 as tetrameric with respect to Ang-1-FD. An angiopoietin-2 version was also constructed and these two molecules were designated Ang-1-FD-Fc-FD and Ang-2-FD-Fc-FD, respectively.

For either type of construct, similar properties were observed: unlike
15 dimeric Ang-1-FD-Fc, which fails to activate Tie-2 in endothelial cells, both Ang-1-FD-FD-Fc and Ang-1-FD-Fc-FD could readily activate Tie-2 in endothelial cells, with a potency comparable to that of the native ligand. Also, like native angiopoietin-2, Ang-2-FD-Fc-FD could antagonize angiopoietin-1 activity with a potency that is comparable to that of native
20 angiopoietin-2, and with much greater potency than the marginally antagonistic activity of the Ang-2-FD-Fc dimer.

Construction of mutant angiopoietin nucleic acid molecules.

25 All of the following nucleic acid molecules were constructed by standard recombinant DNA techniques (See e.g., Molecular Cloning, A Laboratory Manual (Sambrook, et al., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY), Current Protocols in Molecular Biology (Eds. Ausubel, et al., Greene Publ. Assoc., Wiley-Interscience, NY), sequence-verified by standard
30 techniques using an ABI 373A DNA sequencer and Taq Dideoxy Terminator Cycle Sequencing Kit (Applied Biosystems, Inc., Foster City, CA), and subcloned into the mammalian expression vector pMT21 (Genetics

Institute, Inc.) with a Kozak sequence (Kozak, M., 1987, Nucleic Acids Res. 15:8125-8148) at the 5' end to promote protein translation. The bridging sequences described *infra* were introduced to provide convenient restriction sites and to give flexibility to the junctions between the domains, but there is no indication that there is a very critical nature to these bridging sequences (though varying the length of the linker in some of these constructs led to some variation in the amount of protein produced).

Example 1: Construction of the Ang-1-FD-FD-Fc, Ang-2-FD-FD-Fc, Ang-1-FD-Fc-FD, and Ang-2-FD-Fc-FD nucleic acid molecules.

Ang-1-FD-FD-Fc: Ang-1-FD-FD-Fc consists of a trypsin signal sequence at its amino terminus to allow for secretion (bases 1-45 of Figure 1A) followed by the angiopoietin-1 fibrinogen domain (FD) (bases 46-690 of Figure 1A-Figure 1B), a short bridging sequence consisting of the amino acids Gly-Pro Ala-Pro (bases 691-702 of Figure 1B), a second angiopoietin-1 FD (bases 703-1750 of Figure 1B-Figure 1D), another bridging sequence consisting of the amino acids Gly-Pro-Gly (bases 1351-1359 of Figure 1D), and the coding sequence for the Fc portion of human IgG1 (bases 1360-2058 of Figure 1D-Figure 1E).

Ang-2-FD-FD-Fc: The Ang-2-FD-FD-Fc nucleic acid molecule was similarly constructed. It consists of a trypsin signal sequence (bases 1-45 of Figure 2A), an angiopoietin-2 FD (bases 46-690 of Figure 2A- Figure 2B), a bridging amino acid sequence Gly-Gly-Pro-Ala-Pro (bases 691-705 of Figure 2B), a second angiopoietin-2 FD (bases 706-1353 of Figure 2B-Figure 2D), another bridging amino acid sequence Gly-Pro-Gly (bases 1354-1362 of Figure 2D), and the coding sequence for the Fc portion of human IgG1 (bases 1363-2061 of Figure 2D-Figure 2E).

Ang-1-FD-Fc-FD: The Ang-1-FD-Fc-FD consists of a trypsin signal sequence (bases 1-45 of Figure 3A), an angiopoietin-1 FD (bases 46-690 of Figure 3A-3B), the bridging amino acid sequence Gly-Pro-Gly (bases 691-699 of Figure

3B), the coding sequence for the Fc portion of human IgG1 (bases 700-1395 of Figure 3B-3D), another bridging amino acid sequence Gly-Gly-Gly-Gly-Ser-Gly-Ala-Pro (bases 1396-1419 of Figure 3D), and a second angiopoietin-1 FD (bases 1420-2067 of Figure 3D-Figure 3E).

5

Ang-2-FD-Fc-FD: The Ang-2-FD-Fc-FD nucleic acid molecule consists of a trypsin signal sequence (bases 1-45 of Figure 4A), an angiopoietin-2 FD domain (bases 46-690 of Figure 4A-Figure 4B), the bridging amino acid sequence Gly-Gly-Pro-Gly (bases 691-702 of Figure 4B), the coding sequence
10 for the Fc portion of human IgG1 (bases 703-1398 of Figure 4B- Figure 4D), the bridging amino acid sequence Gly-Gly-Gly-Gly-Ser-Gly-Ala-Pro (bases 1399-1422 of Figure 4D), and a second angiopoietin-2 FD (bases 1423-2067 of Figure 4D-Figure 4E).

15 **Example 2: Characterization of Ang-1 FD-Fc-FD protein.**

Molecular Weight Analysis: The predicted molecular weight for Ang-1-FD-Fc-FD protein was determined using the MacVector Program (Kodak, Scientific Imaging Systems, New Haven, CT) The monomeric form (with
20 respect to the Fc) has a predicted weight of 76,349. In addition, there are three predicted N-linked glycosylation sites, approximately 2500 MW/site, that could potentially increase the molecular weight of the monomeric protein to 83,849. Due to the interaction between the Fc components of adjacent molecules, the protein actually exists as a dimer with a predicted
25 molecular weight, including possible N-linked glycosylation, of 167,698. Subsequent SDS PAGE analyses of COS cell-derived protein described *infra* confirmed these approximate molecular weights, with a band running at about 210kD under non-reducing conditions and a band running at about 85kD under reducing conditions (Figure 5). Light scatter analysis was
30 performed to further confirm the molecular weight and, more importantly, determine whether or not the protein was a homogeneous species. Light scattering is a function of mass and concentration of a macromolecule. To

determine molecular weight, the protein sample was injected onto a gel filtration column and the effluent was monitored with an on line light scattering detector and a refractive index and/or a UV detector. The light scattering detector is a MiniDawn laser light scattering detector was from
5 Wyatt Technology Corporation (Santa Barbara, CA). This instrument measures static light at three different angles. The on line refractive index detector or UV detector serve to measure protein concentration. Astra 4.7 Software (Wyatt Technology Corporation, Santa Barbara, CA) was used to calculate the protein concentration based on either dn/dc (dn = change of
10 refractive index; dc = concentration) or the extinction coefficient of the protein. The molecular weight of protein is then calculated from the angular dependence of light scattering. Figure 6 shows the results of this analysis using COS cell-derived protein. The molecular weight of the dimeric protein appears to be approximately 200kD and presence of a single
15 peak implies that the protein solution is, in fact, homogenous.

Expression Level in COS Cells: COS cell supernatant containing recombinant Ang-1-FD-Fc-FD protein was generated by transiently transfecting COS cells with the Ang1-FD-Fc-FD DNA construct described
20 *supra*. All transfections were performed using standard techniques known in the art. The COS cell supernatant was analyzed using Biacore technology (Pharmacia, Inc.) to quantitate the amount of Ang-1-FD-Fc-FD protein present in the supernatant. This analysis resulted in an RU value of 765, which is equivalent to 0.9mg of recombinant protein/liter of COS cell
25 supernatant. These values represent very high levels of expression.

Purification of COS Supernatants: Because the Ang-1-FD-Fc-FD protein contains an Fc domain, purification is relatively simple and straight forward using standard Protein A column chromatography (Pharmacia,
30 Inc.) followed by standard size exclusion chromatography (Pharmacia, Inc.). In fact, the relative ease of purification of the Ang-1-FD-Fc-FD protein gives it a distinct advantage over the parent protein, angiopoietin-1, from which

it is derived, and the mutant version of angiopoietin-1 called Ang1* that consists of the N-terminal of angiopoietin-2 fused to the coiled-coil domain and fibrinogen domain of angiopoietin-1 and that has a Cys to Ser mutation at amino acid 245. (See PCT International Application entitled "Modified
5 TIE-2 Receptor Ligands," published as WO 98/05779 on 12 February 1998 in the name of Regeneron Pharmaceuticals, Inc., especially Figure 27, which is hereby incorporated by reference).

Both angiopoietin-1 and Ang1* require extensive, expensive and labor-
10 intensive purification schemes that result in relatively poor yields of recombinant protein. The need for cost-effective, simple purification schemes for biologicals intended for clinical use can not be over-emphasized.

15 The COS cell supernatant was purified as described *supra* and yielded approximately 1 mg of purified Ang-1-FD-Fc-FD protein that was used in the studies described *infra* to further characterize the protein.

N-terminal sequencing of COS cell-derived Ang-1-FD-Fc-FD protein:

20 Purified Ang-1-FD-Fc-FD protein was subjected to standard N-terminal sequence analysis to determine if any truncated species of the protein were being generated. This was of concern because the mutant molecule, Ang1*, has a history of containing between 10-20% N-terminally truncated species. The analysis revealed only one N-terminal sequence, Arg-Asp-X-Ala-Asp,
25 wherein X is Cys. This sequence can be found at amino acids 16-20 of Figure 3A, and immediately follows the protein's signal sequence corresponding to amino acids 1-15 Figure 3A.

Receptor binding analysis of COS cell-derived Ang-1-FD-Fc-FD: Previous

30 studies have determined that the fibrinogen domain (FD) of the angiopoietin molecules is necessary for ligand/receptor interaction. Furthermore, in order for high affinity binding to the Tie-2 receptor to

occur, native angiopoietin-1, angiopoietin-2, and the mutant Ang1* must form at least tetrameric, and possibly higher order, multimers. To determine whether the COS cell-derived Ang-1-FD-Fc-FD protein, which is tetrameric with respect to the FD domain, could bind to Tie-2 with high affinity, standard Biacore analysis was performed. Briefly, Tie-2-Fc receptor body protein, which is a fusion protein comprising the ectodomain of Tie-2 fused to the Fc domain of human IgG1, was immobilized on a Biacore chip. Ang-1-FD-Fc-FD-containing solution was passed over the chip and binding between Tie-2 ectodomain and Ang-1-FD-Fc-FD was allowed to occur. The binding step was followed by a 0.5 M NaCl high salt wash. The high salt wash was not able to disrupt the interaction between the Ang-1-FD-Fc-FD protein and the Tie-2 receptor ectodomain, implying that there is a strong interaction between the mutant ligand and receptor. This result is consistent with earlier Biacore results in which both Ang-1-FD-Fc-FD parent molecule, angiopoietin-1 and the mutant Ang1* molecule, have been shown to interact strongly with the Tie-2-Fc receptor and that this interaction is not disrupted by high salt. In contrast, several mutant molecules derived from the parent angiopoietin-1 molecule are readily dissociated from the Tie-2-Fc receptor when treated with high salt. The mutant molecules, designated Ang-1/FD (a monomer with respect to the FD), Ang-1/FD-Fc (also a monomer with respect to the FD, but which is able to form a dimer due to the presence of the Fc domain), and Ang-1/C/FD (a monomer with respect to the FD, but which also contains the coiled-coil domain of angiopoietin-1), do not exist in multimeric forms sufficient for high affinity binding to the Tie-2 receptor.

Example 3: Characterization of COS cell-derived Ang-2-FD-Fc-FD protein.

Molecular Weight Analysis: As described for Ang-1-FD-Fc-FD *supra*, the predicted molecular weight for Ang-2-FD-Fc-FD protein was determined using the MacVector Program (Kodak, Scientific Imaging Systems, New Haven, CT) The monomeric form of Ang-2-FD-Fc-FD has a predicted

- weight of 76,052, with three predicted N-linked glycosylation sites that could potentially increase the molecular weight of the monomeric protein to 83,552. Like Ang-1-FD-Fc-FD, the protein exists as a dimer with a predicted molecular weight, including possible N-linked glycosylation, of 167,104.
- 5 SDS PAGE analyses of COS cell-derived protein confirmed these approximate molecular weights, with a band running at about 200kD under non-reducing conditions and a band running at about 88kD under reducing conditions (Figure 7). Light scatter analysis confirmed the molecular weight (171kD) and revealed that the Ang-2-FD-Fc-FD protein, like Ang-1-FD-Fc-
- 10 FD, exists as a homogeneous species (Figure 8).

Expression Level in COS Cells: COS cell supernatant containing recombinant Ang-2-FD-Fc-FD protein was generated by transiently transfecting COS cells with the Ang-2-FD-Fc-FD DNA construct described

15 *supra*. The COS cell supernatant was analyzed by Biacore to quantitate the amount of Ang-2-FD-Fc-FD protein present in the supernatant. This analysis resulted in an RU value of 606, which is equivalent to 0.7mg of recombinant protein/liter of COS cell supernatant. These values represent relatively high levels of expression.

20

Purification of COS Supernatants: As with Ang-1-FD-Fc-FD, Ang-2-FD-Fc-FD protein contains an Fc domain, so purification is relatively simple and straight forward using standard Protein A column chromatography followed by standard size exclusion chromatography. The COS cell

25 supernatant was purified as described for Ang-1-FD-Fc-FD *supra* and yielded approximately 2 mg of purified Ang-2-FD-Fc-FD protein that was used in the studies described *infra* to further characterize this protein.

N-terminal sequencing: Purified COS cell-derived Ang-2-FD-Fc-FD protein

30 was subjected to standard N-terminal sequence analysis to determine if any truncated species of the protein were being generated. The analysis revealed only one N-terminal sequence, Arg-Asp-X-Ala-Glu, wherein X is Cys. This

sequence can be found at amino acids 16-20 of Figure 4A, and immediately follows the protein's signal sequence corresponding to amino acids 1-15 of Figure 4A.

- 5 **Receptor binding analysis of COS cell-derived protein:** To determine whether the COS cell-derived Ang-2-FD-Fc-FD protein could bind to the Tie-2 receptor, standard Biacore analysis was performed as described for Ang-1-FD-Fc-FD *supra*. As with Ang-1-FD-Fc-FD, a high salt wash was not able to disrupt the interaction between the Ang-2-FD-Fc-FD protein and the
10 Tie-2-Fc receptor, again implying that there is a strong interaction between mutant ligand and receptor.

Example 4: Effects of COS cell-derived Ang-1-FD-Fc-FD and Ang-2-FD-Fc-FD on Tie-2 receptor phosphorylation in EAhy926 cells.

- 15 Because Ang-1-FD-Fc-FD is a mutant molecule derived from the agonist angiopoietin-1 and Ang-2-FD-Fc-FD is a mutant molecule derived from the antagonist angiopoietin-2, we wanted to determine whether or not these two mutant molecules would retain the activity associated with the parent
20 molecule from which it was derived.

- Assay system:** All of the experiments described *infra* utilized the cell line EAhy926 (Edgell, C. J., et al., (1983) Proc. Natl. Acad. Sci. USA 80:3734-3737) and standard phosphorylation assays and reagents familiar to those of skill
25 in the art.

- (A) Ang1*-mediated vs. Ang-1-FD-Fc-FD-mediated Tie-2 receptor phosphorylation in EAhy926 cells:** EAhy926 cells were stimulated with either 0.1 µg/ml, 0.2 µg/ml, or 0.8 µg/ml Ang1* or Ang-1-FD-Fc-FD protein.
30 A standard phosphorylation assay revealed that Ang-1-FD-Fc-FD was equivalent to Ang1* in its ability to stimulate phosphorylation of the Tie-2 receptor in EAhy926 cells (Figure 9).

(B) Ability of Ang-2-FD-Fc-FD to block Ang1*-mediated Tie-2 receptor phosphorylation in EAhy926 cells: EAhy926 cells were treated with 0.4 µg/ml of the Tie-2 agonist Ang1* and 1 µg/ml, 2 µg/ml, 4 µg/ml, 6 µg/ml, or 8 µg/ml of Ang-2-FD-Fc-FD. As shown in Figure 10, Ang-2-FD-Fc-FD is able to block Ang1* stimulation of the Tie-2 receptor when it is present in at least a 10-15 fold molar excess of Ang1*.

(C) Ability of angiopoietin-2 to block Ang1*-mediated Tie-2 receptor phosphorylation in EAhy926 cells: To compare the blocking effects of the naturally occurring antagonist angiopoietin-2 with that of Ang-2-FD-Fc-FD, the same experiment described in (B) *supra* was performed, substituting angiopoietin-2 for Ang-2-FD-Fc-FD. The results of this experiment are shown in Figure 11. At a 20 fold molar excess, the angiopoietin-2 has not reduced the phosphorylation level to 50%. This result, coupled with the results described in (B) *supra* implies that Ang-2-FD-Fc-FD is a more potent inhibitor of Ang1*-mediated Tie-2 receptor phosphorylation than angiopoietin-2.

(D) Ability of Ang-2-FD-Fc-FD to block angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in EAhy926 cells: EAhy926 cells were treated with 0.2 µg/ml of the naturally occurring Tie-2 agonist angiopoietin-1 and 1 µg/ml, 2 µg/ml, 4 µg/ml, 6 µg/ml, or 8 µg/ml of Ang-2-FD-Fc-FD. The results of this experiment, shown in Figure 12, show that while there is a trend toward blocking angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in these cells, Ang-2-FD-Fc-FD seems to be more effective at blocking Ang1*-mediated phosphorylation of Tie-2, as shown in Figure 10 and described in (B) *supra*.

(E) Ability of angiopoietin-2 to block angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in EAhy926 cells: EAhy926 cells were treated with 0.2 µg/ml of the angiopoietin-1 and 1 µg/ml, 2 µg/ml, 4 µg/ml,

6 µg/ ml, or 8 µg/ml of angiopoietin-2. The results of this experiment, shown in Figure 13, show that there is a trend toward blocking angiopoietin-1-mediated phosphorylation of the Tie-2 receptor in these cells, but, like Ang-2-FD-Fc-FD, angiopoietin-2 seems to be more effective at blocking Ang1*-mediated phosphorylation of Tie-2, as shown in Figure 11 and described in (C) *supra*.

Example 5: Construction of Ang-1-FD-Fc-FD CHO cell expression vector pRG763/Ang-1-FD-Fc-FD.

10

The pRG763/Ang-1-FD-Fc-FD CHO cell expression vector was constructed by isolating from the plasmid pCDNA3.1/Ang1-FD-Fc-FD a 2115 base pair EcoRI - NotI fragment containing Ang1-FD-Fc-FD and ligating this fragment into pRG763 vector digested with EcoRI and NotI. A large scale (2L) culture of E. coli DH10B cells carrying the pRG763/Ang-1-FD-Fc-FD plasmid was grown overnight in TB + ampicillin and the plasmid DNA was extracted using a Promega Wizard Plus Maxiprep kit, following the manufacturer's protocol. The concentration of the purified plasmid DNA was determined in a UV spectrophotometer and fluorometer. The plasmid DNA was verified by digestion of aliquots with NcoI and HincII restriction enzymes. All restriction enzyme digest fragments corresponded to the predicted sizes in a 1% agarose gel.

20

Example 6: Expression of Ang-1-FD-Fc-FD in CHO cells.

25

Forty 15 cm petri plates were seeded with CHO-K1/E1A cells at a density of 4×10^6 cells/plate. Plating media was Gibco Ham's F-12 w/10% Hyclone Fetal Bovine Serum (FBS) + penicillin/streptomycin and supplemented with glutamine. The following day each plate was transfected with 6 µg of pRG763/Ang-1-FD-Fc-FD using Gibco Optimem and Gibco Lipofectamine in 12 ml volume, following the manufacturer's protocol. Four hours after adding the transfection mix to the cells 12 ml/plate of Optimem w/ 10%

30

FBS was added. Plates were incubated at 37°C in a 5% CO₂ incubator overnight. The following day the media was removed from each plate and 25 ml expression media (Gibco CHO-S-SFM II w/ glutamine + 1mM sodium butyrate) was added. The plates were incubated at 37°C for 3 days. After 3
5 days of incubation the media was aspirated from each plate and centrifuged at 400 rpm in a swinging bucket rotor to pellet cells. The supernatant was decanted into sterile 1L bottles and purified as described *infra*.

Example 7: Construction of Ang-2-FD-Fc-FD CHO cell expression vector
10 **pRG763/Ang-2-FD-Fc-FD.**

The plasmid pRG763/Ang-2-FD-Fc-FD was constructed by isolating from the plasmid pCDNA3.1/Ang-2-FD-Fc-FD a 2097 base pair EcoRI - NotI fragment containing Ang-2-FD-Fc-FD and ligating this fragment into the pRG763
15 vector digested with EcoRI and NotI. A large scale (1L) culture of E. coli DH10B cells carrying the pRG763/Ang-2-FD-Fc-FD plasmid was grown overnight in TB + ampicillin and the plasmid DNA was extracted using a Promega Wizard Plus Maxiprep kit, following the manufacturer's protocol. The concentration of the purified plasmid DNA was determined in a UV
20 spectrophotometer and fluorometer. The plasmid DNA was also verified by digestion of plasmid DNA with NcoI and Ppu10I restriction enzymes. All restriction enzyme digest fragments corresponded to the predicted sizes in a 1% agarose gel.

25 **Example 8: Expression of Ang-2-FD-Fc-FD in CHO cells.**

Forty 15 cm petri plates were seeded with CHO-K1/E1A cells at a density of 4 x 10⁶ cells/plate. Plating media was Gibco Ham's F-12 w/10% Hyclone Fetal Bovine Serum (FBS) + penicillin/streptomycin and supplemented with
30 glutamine. The following day each plate was transfected with 6 µg of pRG763/Ang-2-FD-Fc-FD using Gibco OptiMem and Gibco Lipofectamine in 12 ml volume, following the manufacturer's protocol. Four hours after

adding the transfection mix to the cells 12 ml/plate of Optimem w/ 10% FBS was added. Plates were incubated at 37°C in a 5% CO₂ incubator overnight. The following day the media was removed from each plate and 25 ml expression media (Gibco CHO-S-SFM II w/ glutamine + 1mM sodium butyrate) was added. The plates were incubated at 37°C for 3 days. After 3 days of incubation the media was aspirated from each plate and centrifuged at 400 rpm in a swinging bucket rotor to pellet cells. The supernatant was decanted into sterile 1L bottles purified as described infra.

10 **Example 9: Characterization of stable CHO clone-derived Ang-1-FD-Fc-FD protein.**

Molecular Weight Analysis: The predicted molecular weight for stable CHO clone-derived Ang-1-FD-Fc-FD protein was determined using the MacVector Program (Kodak, Scientific Imaging Systems, New Haven, CT) The monomeric form (with respect to the Fc) has a predicted weight of 76,349. In addition, there are three predicted N-linked glycosylation sites, approximately 2500 MW/site, that could potentially increase the molecular weight of the monomeric protein to 83,849. Due to the interaction between the Fc components of adjacent molecules, the protein actually exists as a dimer with a predicted molecular weight, including possible N-linked glycosylation, of 167,698. Subsequent SDS PAGE analyses confirmed these approximate molecular weights, with a band running at about 210kD under non-reducing conditions and a band running at about 85kD under reducing conditions. Light scatter analysis was performed to further confirm the molecular weight and, more importantly, determine whether or not the protein was a homogeneous species. Light scattering is a function of mass and concentration of a macromolecule. To determine molecular weight, the protein sample was injected onto a gel filtration column and the effluent was monitored with an on line light scattering detector and a refractive index and/or a UV detector. The light scattering detector is a MiniDawn laser light scattering detector was from Wyatt Technology

Corporation (Santa Barbara, CA). This instrument measures static light at three different angles. The on line refractive index detector or UV detector serve to measure protein concentration. Astra 4.7 Software (Wyatt Technology Corporation, Santa Barbara, CA) was used to calculate the
5 protein concentration based on either dn/dc (dn = change of refractive index; dc = concentration) or the extinction coefficient of the protein. The molecular weight of protein is then calculated from the angular dependence of light scattering. The results of this analysis show that the dimeric protein appears to be approximately 173.9kD and the presence of a single peak
10 implies that the protein solution is homogenous.

Expression level of Ang-1-FD-Fc-FD in stable CHO clones: CHO cell supernatant containing recombinant Ang-1-FD-Fc-FD protein was generated by stably transfecting CHO cells with the Ang-1-FD-Fc-FD DNA construct
15 described *supra*. The CHO cell supernatant was analyzed by standard ELISA using an anti-human IgG antibody as a capture antibody and an anti-human IgG antibody conjugated to alkaline phosphatase as a reporter antibody to quantitate the amount of Ang-1-FD-Fc-FD protein present in the supernatant. This analysis revealed expression levels of 2-3 pg/cell/day.

20
Purification of Ang-1-FD-Fc-FD protein derived from stable CHO clone supernatants: Because the Ang-1-FD-Fc-FD protein contains an Fc domain, purification is relatively simple and straight forward using standard Protein A column chromatography (Pharmacia, Inc.) followed by standard size
25 exclusion chromatography (Pharmacia, Inc.). The CHO cell supernatant was purified as described *supra* and the purified ANG-1-FD-Fc-FD protein was used in the studies described *infra* to further characterize the protein.

N-terminal sequencing of stable CHO clone-derived Ang-1-FD-Fc-FD protein: Purified Ang-1-FD-Fc-FD protein was subjected to standard N-
30 terminal sequence analysis to determine if any truncated species of the protein were being generated. The analysis revealed only one N-terminal

sequence, Arg-Asp-X-Ala-Asp, wherein X is Cys. This sequence can be found at amino acids 16-20 of Figure 3A, and immediately follows the protein's signal sequence corresponding to amino acids 1-15 Figure 3A.

5 **Example 10: Characterization of stable CHO clone-derived Ang-2-FD-Fc-FD protein.**

Molecular Weight Analysis: As described for stable CHO clone-derived Ang-1-FD-Fc-FD *supra*, the predicted molecular weight for stable CHO
10 clone-derived Ang-2-FD-Fc-FD protein was determined using the MacVector Program (Kodak, Scientific Imaging Systems, New Haven, CT). The monomeric form of Ang-2-FD-Fc-FD has a predicted weight of 76,052, with three predicted N-linked glycosylation sites that could potentially increase the molecular weight of the monomeric protein to 83,552. Like
15 Ang-1-FD-Fc-FD, the protein exists as a dimer with a predicted molecular weight, including possible N-linked glycosylation, of 167,104. SDS PAGE analyses confirmed these approximate molecular weights, with a band running at about 200kD under non-reducing conditions and a band running at about 85kD under reducing conditions. Light scatter analysis confirmed
20 the molecular weight (176.6kD) and revealed that the stable CHO clone-derived Ang-2-FD-Fc-FD protein, like stable CHO clone-derived Ang-1-FD-Fc-FD, exists as a homogeneous species.

Expression level of Ang-2-FD-Fc-FD derived from stable CHO clones: CHO
25 cell supernatant containing recombinant Ang-2-FD-Fc-FD protein was generated by stably transfecting CHO cells with the Ang-2-FD-Fc-FD DNA construct described *supra*. The CHO cell supernatant was analyzed by standard ELISA using an anti-human IgG antibody as a capture antibody and an anti-human IgG antibody conjugated to alkaline phosphatase as a
30 reporter antibody to quantitate the amount of Ang-2-FD-Fc-FD protein present in the supernatant. This analysis revealed expression levels of approximately 1-2 pg/cell/day.

Purification of stable CHO clone-derived Ang-2-FD-Fc-FD from cell

supernatants: As with Ang-1-FD-Fc-FD, Ang-2-FD-Fc-FD protein contains an Fc domain, so purification is relatively simple and straight forward using standard Protein A column chromatography followed by standard size exclusion chromatography. The CHO cell supernatant was purified as described for stable CHO clone-derived Ang-1-FD-Fc-FD *supra* and was used in the studies described *infra* to further characterize this protein.

N-terminal sequencing of stable CHO clone-derived Ang-2-FD-Fc-FD

protein: Purified stable CHO clone-derived Ang-2-FD-Fc-FD protein was subjected to standard N-terminal sequence analysis to determine if any truncated species of the protein were being generated. The analysis revealed only one N-terminal sequence, Asp-X-Ala-Glu-Val, wherein X is Cys. This sequence can be found at amino acids 17-21 of Figure 4A, and immediately follows the protein's signal sequence corresponding to amino acids 1-15 of Figure 4A.

Example 11: Effects of stable CHO clone-derived Ang-1-FD-Fc-FD and Ang-2-FD-Fc-FD on Tie-2 receptor phosphorylation in EAhy926 cells.

Assay system: All of the experiments described *infra* utilized the cell line EAhy926 (Edgell, C. J., et al., (1983) Proc. Natl. Acad. Sci. USA 80:3734-3737) and standard phosphorylation assays and reagents familiar to those of skill in the art.

(A) Ang1*-mediated vs. stable CHO clone-derived Ang-1-FD-Fc-FD-mediated Tie-2 receptor phosphorylation in EAhy926 cells: EAhy926 cells were stimulated with 0.4 µg/ml Ang1* or 0.2 µg/ml or 0.4 µg/ml stable CHO clone-derived Ang-1-FD-Fc-FD protein. A standard phosphorylation assay revealed that or stable CHO clone-derived Ang-1-FD-Fc-FD was equivalent to Ang1* in its ability to stimulate phosphorylation of the Tie-2 receptor in EAhy926 cells (Figure 17).

(B) Ability of stable CHO clone-derived Ang-2-FD-Fc-FD to block stable CHO clone-derived Ang-1-FD-Fc-FD-mediated Tie-2 receptor phosphorylation in EAhy926 cells:

EAhy926 cells were treated with 0.2 µg/ml of the Tie-2 agonist Ang-1-FD-Fc-FD and 2 µg/ml, 4 µg/ml, 8 µg/ml or 16 µg/ml of stable CHO clone-derived Ang-2-FD-Fc-FD. As shown in Figure 18, Ang-2-FD-Fc-FD is able to block stable CHO clone-derived Ang-1-FD-Fc-FD stimulation of the Tie-2 receptor when it is present in at least a 40 fold molar excess of stable CHO clone-derived Ang-1-FD-Fc-FD.

Ephrin ligands:

In previous experiments (Davis et al., 1994, Science, 266:816-819; Gale et al., 1996, Neuron 17:9-19, Gale and Yancopoulos, 1997, Cell Tissue Research 290:227-241), soluble, unclustered Ephrin-B1-Fc and Ephrin-B2-Fc, which dimerize at their respective Fc domains and therefore are dimeric with respect to either the Ephrin-B1 or Ephrin-B2 ectodomain, failed to induce EphB2 receptor phosphorylation. However, when either molecule was multimerized by pre-clustering with an anti-Fc antibody, they became potent agonists for the EphB2 receptor, as judged by tyrosine phosphorylation of the EphB2 receptor in a COS cell reporter assay. Because multimerization of both Ephrin-B1 and Ephrin-B2 appears to be necessary for induction of receptor phosphorylation, we theorized that a molecule that contained tandem repeats of either Ephrin-B1 or Ephrin-B2 ectodomains fused to an Fc domain, which would be dimeric with respect to the Fc domain but which would be tetrameric with respect to Ephrin ectodomains, might be sufficiently clustered to induce receptor phosphorylation. To test this hypothesis, the following DNA constructs were constructed, recombinant proteins produced, and reporter assays performed.

Construction of tandem Ephrin ectod main/Fc domain nucleic acid molecules.

All of the following nucleic acid molecules were constructed by standard recombinant DNA techniques (See e.g., Molecular Cloning, A Laboratory Manual (Sambrook, et al., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY), Current Protocols in Molecular Biology (Eds. Ausubel, et al., Greene Publ. Assoc., Wiley-Interscience, NY), sequence-verified by standard techniques using an ABI 373A DNA sequencer and Taq Dideoxy Terminator Cycle Sequencing Kit (Applied Biosystems, Inc., Foster City, CA), and subcloned into either the mammalian expression pJFE14 (Ephrin-B1-Ephrin-B1-Fc) or pMT21 (Ephrin-B2-Ephrin-B2-Fc), each with a Kozak sequence (Kozak, M., 1987, Nucleic Acids Res. 15:8125-8148) at the 5' end to promote protein translation. The bridging sequences described *infra* were introduced to provide convenient restriction sites and to give flexibility to the junctions between the domains, but there is no indication that there is a very critical nature to these bridging sequences (though varying the length of the linker in some of these constructs led to some variation in the amount of protein produced).

20

Example 12: Construction of Ephrin-B1-Ephrin-B1-Fc and Ephrin-B2-Ephrin-B2-Fc nucleic acid molecules.

(A) Ephrin-B1-Ephrin-B1-Fc: The Ephrin-B1-Ephrin-B1-Fc DNA molecule consists of the coding sequence of the ectodomain of Ephrin-B1 (Davis et al., *ibid.*), which corresponds to nucleotides 1-711 of Figure 14A-Figure 14B, followed by a bridging sequence consisting of the amino acids Gly-Pro-Gly (nucleotides 712-720 of Figure 14B), followed by a second copy of the ectodomain of Ephrin-B1 (corresponding to nucleotides 721-1344 of Figure 14B-Figure 14D), except that in this copy of the Ephrin-B1 ectodomain the signal sequence has been removed. This second copy is followed by a second Gly-Pro-Gly amino acid bridge (nucleotides 1345-1353 of Figure 14D),

30

followed by the coding sequence for the Fc portion of human IgG1 (nucleotides 1354-2049 of Figure 14D-Figure 14E).

(B) Ephrin-B2-Ephrin-B2-Fc: The Ephrin-B2-Ephrin-B2-Fc DNA molecule consists of the coding sequence of the ectodomain of Ephrin-B2 (Bergemann et al., 1995, Mol. Cell Biol. 15:4821-4929), which corresponds to nucleotides 1-675 of Figure 15A-Figure 15B, followed by a bridging sequence consisting of the amino acids Gly-Pro-Gly (nucleotides 676-684 of Figure 15B), followed by a second copy of the ectodomain of Ephrin-B2 (corresponding to nucleotides 685-1270 of Figure 15B-Figure 15D), except that in this copy the signal sequence has been removed. This second copy is followed by a second Gly-Pro-Gly amino acid bridge (nucleotides 1270-1278 of Figure 15D), followed by the coding sequence for the Fc portion of human IgG1 (nucleotides 1279-1977 of Figure 15D-Figure 15E).

As with the angiopoietin nucleic acid molecules described *supra*, the bridging sequences were introduced to provide convenient restriction sites and to give flexibility to the junctions between the domains.

Example 13: Expression of tandem Ephrin recombinant proteins in COS cells.

COS cells were transiently transfected with either the Ephrin-B1-Ephrin-B1-Fc or Ephrin-B2-Ephrin-B2-Fc nucleic acid molecules described *supra* using standard transfection techniques known in the art. Two days subsequent to transfection, the growth medium (DMEM supplemented with 100 U/ml penicillin, 100 µg/ml streptomycin, 2 mM glutamine, and 10% calf serum) was aspirated and replaced with serum-free medium (DMEM supplemented with 100 U/ml penicillin, 100 µg/ml streptomycin, 2 mM glutamine). Cell were grown for an additional three days and then the serum-free medium containing the recombinant proteins was collected. Recombinant protein concentration was determined by performing dot blots and comparing the

signal obtained to a standard curve. Once approximate protein concentrations were determined, the Ephrin-B1-Ephrin-B1-Fc and Ephrin-B2-Ephrin-B2-Fc recombinant proteins were used in the cell reporter assays described *infra*.

5

Example 14: Characterization of the COS cell-derived tandem Ephrin ectodomain/Fc domain recombinant proteins.

Reporter Assay: COS cells, which endogenously express the Eph family
10 receptor EphB2 (Gale et al., 1996, Neuron 17:9-19), were used in reporter assays to evaluate the ability of Ephrin-B1-Ephrin-B1-Fc and Ephrin-B2-Ephrin-B2-Fc to induce receptor phosphorylation. The assays were performed as previously described (Davis et al., *ibid.*; Gale et al., *ibid.*). Briefly, COS cells were grown to 80-90% confluency in standard growth
15 medium described *supra*. After growth, the medium was aspirated, and replaced with serum-free medium (described *supra*) for 1-2 hours prior to treatment with either Ephrin-B1-Ephrin-B1-Fc or Ephrin-B2-Ephrin-B2-Fc recombinant protein. The cells were stimulated with 500 ng/ml Ephrin-B1-Ephrin-B1-Fc or Ephrin-B2-Ephrin-B2-Fc for 30 minutes at 37°C, with or
20 without affinity purified human IgG1 Fc-specific goat anti-human antibody (Jackson Immunoresearch, West Grove, PA) at a final concentration of 17 µg/ml. This antibody is capable of clustering the Fc tagged fusion. Subsequent to treatment, the COS cells were harvested and cell lysates were prepared as described in Davis, et al. and Gale, et al., *supra*. The EphB2
25 receptor protein was immunoprecipitated from the cell lysates using an anti-EphB2 antisera (Henkemeyer et al., 1994, Oncogene 9:1001-1014). Immunoprecipitates were resolved by standard SDS PAGE and transferred to PVDF membranes (Millipore) for western blot analysis. The membranes were probed with either anti-phosphotyrosine antibody 4G10 (Upstate
30 Biotechnology Institute, Lake Placid, NY) or anti-EphB2 antibodies (Henkemeyer, et al., *ibid.*) to determine the extent of EphB2

phosphorylation and the relative quantities of EphB2 in the experimental conditions described *supra*.

Results: Both Ephrin-B1-Ephrin-B1-Fc and Ephrin-B2-Ephrin-B2-Fc were shown to be approximately as active as anti-Fc antibody-clustered Ephrin-B1-Fc in their ability to induce EphB2 receptor phosphorylation in the COS cell reporter assay. Furthermore, if either of the proteins were further clustered with the goat anti-human Fc antibody, they became even more potent in their ability to induce EphB2 receptor phosphorylation. Figure 16 shows the results of this phosphorylation assay.

Example 15: Construction of Ephrin-B2-Ephrin-B2-Fc CHO expression vector.

The Ephrin-B2-Ephrin-B2-Fc DNA molecule consists of the coding sequence of the ectodomain of Ephrin-B2 (Bergemann et al., 1995, Mol. Cell Biol. 15:4821-4929), which corresponds to nucleotides 1-675 of Figure 15A-Figure 15B, followed by a bridging sequence consisting of the amino acids Gly-Pro-Gly (nucleotides 676-684 of Figure 15B), followed by a second copy of the ectodomain of Ephrin-B2 (corresponding to nucleotides 685-1270 of Figure 15B-Figure 15D), except that in this copy the signal sequence has been removed. This second copy is followed by a second Gly-Pro-Gly amino acid bridge (nucleotides 1270-1278 of Figure 15D), followed by the coding sequence for the Fc portion of human IgG1 (nucleotides 1279-1977 of Figure 15D-Figure 15E). This molecule was subcloned into the HindIII and NotI polylinker sites in the expression vector pRG763 and was designated pRG763-m(Ephrin-B2)₂-Fc. As with the angiopoietin nucleic acid molecules described *supra*, the bridging sequences were introduced to provide convenient restriction sites and to give flexibility to the junctions between the domains.

Example 16: Expression of Ephrin-B2-Ephrin-B2-Fc in CHO-K1 (E1A) cells.

Forty 15 cm petri plates were seeded with CHO-K1/E1A cells at a density of 4×10^6 cells/plate. Plating media was Gibco Ham's F-12 w/10% Hyclone Fetal Bovine Serum (FBS) + penicillin/streptomycin and supplemented with glutamine. The following day each plate was transfected with 6 μ g of pRG763-m(Ephrin-B2)2-Fc using Gibco Optimem and Gibco Lipofectamine in 12 ml volume, following the manufacturer's protocol. Four hours after adding the transfection mix to the cells 12 ml/plate of Optimem w/ 10% FBS was added. Plates were incubated at 37°C in a 5% CO₂ incubator overnight. The following day the media was removed from each plate and 25 ml expression media (Gibco CHO-S-SFM II w/ glutamine + 1mM sodium butyrate) was added. The plates were incubated at 37°C for 3 days. After 3 days of incubation the media was aspirated from each plate and centrifuged at 400 rpm in a swinging bucket rotor to pellet cells. The supernatant was decanted into sterile 1L bottles and purified as described *supra*.

WHAT IS CLAIMED IS:

1. A nucleic acid encoding a fusion polypeptide wherein the fusion polypeptide comprises a first subunit comprising at least one copy of the receptor binding domain of a ligand, the first subunit being fused to the N-terminal end of a multimerizing component, said multimerizing component being fused at its C-terminal end to a second subunit comprising at least one copy of the receptor binding domain of a ligand.
2. The nucleic acid of claim 1, wherein the receptor binding domains of the first and second subunits are copies of the receptor binding domain of the same ligand.
3. The nucleic acid of claim 1, wherein the receptor binding domains of the first subunit are copies of the receptor binding domain of a different ligand from the receptor binding domains of the second subunit.
4. The nucleic acid of claim 2, wherein the first and second subunits each have one copy of the receptor binding domain of the ligand.
5. The nucleic acid of claim 3, wherein the first and second subunits each have one copy of the receptor binding domain of the ligand.
6. The nucleic acid of claim 2, wherein the receptor binding domain is the fibrinogen domain of angiopoietin-1 or angiopoietin-2.
7. The nucleic acid of claim 4, wherein the receptor binding domain is the fibrinogen domain of angiopoietin-1 or angiopoietin-2.

8. The nucleic acid of claim 2, wherein the ligand is selected from the group consisting of the EPH family of ligands.
9. The nucleic acid of claim 4, wherein the ligand is selected from the group consisting of the EPH family of ligands.
10. The nucleic acid of claims 1 through 9, wherein the multimerizing component comprises an immunoglobulin derived domain.
11. The nucleic acid molecule of claim 10, wherein the immunoglobulin derived domain is selected from the group consisting of the Fc domain of IgG, the heavy chain of IgG, and the light chain of IgG.
12. A fusion polypeptide encoded by the nucleic acid molecule of claims 1 through 11.
13. A composition comprising a multimer of the fusion polypeptide of claim 12.
14. The composition of claim 13, wherein the multimer is a dimer.
15. A vector which comprises the nucleic acid molecule of claims 1 through 11.
16. An expression vector comprising a nucleic acid molecule of claims 1 through 11, wherein the nucleic acid molecule is operatively linked to an expression control sequence.
17. A host-vector system for the production of a fusion polypeptide which comprises the expression vector of claim 16, in a suitable host cell.

18. The host-vector system of claim 17, wherein the suitable host cell is a bacterial cell, yeast cell, insect cell or mammalian cell.
19. The host-vector system of claim 17, wherein the suitable host cell is E. coli.
20. The host-vector system of claim 17, wherein the suitable host cell is a COS cell.
21. The host-vector system of claim 17, wherein the suitable host cell is a CHO cell.
22. A method of producing a fusion polypeptide which comprises growing cells of the host-vector system of claims 17 through 21, under conditions permitting production of the fusion polypeptide and recovering the polypeptide so produced.
23. A nucleic acid encoding a fusion polypeptide wherein the fusion polypeptide comprises more than one copy of the receptor binding domain of a ligand in tandem, and wherein either the N-terminal or the C-terminal receptor binding domain is also fused to a multimerizing component.
24. The nucleic acid of claim 23, wherein the receptor binding domains are fused contiguously.
25. The nucleic acid of claim 23, wherein the ligand is not a member of the EPH family of ligands.
26. The nucleic acid of claim 24, wherein the ligand is not a member of

the EPH family of ligands.

27. The nucleic acid of claim 23, wherein the receptor binding domain is the fibrinogen domain of angiopoietin-1 or angiopoietin-2.
28. The nucleic acid of claim 24, wherein the receptor binding domain is the fibrinogen domain of angiopoietin-1 or angiopoietin-2.
29. The nucleic acid of claims 23 through 28, wherein the multimerizing component comprises an immunoglobulin derived domain.
30. The nucleic acid molecule of claim 29, wherein the immunoglobulin derived domain is selected from the group consisting of the Fc domain of IgG, the heavy chain of IgG, and the light chain of IgG.
31. A fusion polypeptide encoded by the nucleic acid molecule of claims 23 through 30.
32. A composition comprising a multimer of the fusion polypeptide of claim 31.
33. The composition of claim 32, wherein the multimer is a dimer.
34. A vector which comprises the nucleic acid molecule of claims 23 through 30.
35. An expression vector comprising a nucleic acid molecule of claims 23 through 30, wherein the nucleic acid molecule is operatively linked to an expression control sequence.

36. A host-vector system for the production of a fusion polypeptide which comprises the expression vector of claim 35, in a suitable host cell.
37. The host-vector system of claim 36, wherein the suitable host cell is a bacterial cell, yeast cell, insect cell or mammalian cell.
38. The host-vector system of claim 36, wherein the suitable host cell is E. coli.
39. The host-vector system of claim 36, wherein the suitable host cell is a COS cell.
40. The host-vector system of claim 36, wherein the suitable host cell is a CHO cell.
41. A method of producing a fusion polypeptide which comprises growing cells of the host-vector system of claims 36 through 40, under conditions permitting production of the fusion polypeptide and recovering the polypeptide so produced.

```

      10      20      30      40
*   *   *   *   *   *   *   *
ATG TCT GCA CTT CTG ATC CTA GCT CTT GTT GGA GCT GCA GTT GCT
Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val Ala>
_a_a_a_a_a_TRYPSIN SIGNAL SEQUENCE_a_a_a_a_a>

      50      60      70      80      90
*   *   *   *   *   *   *   *
AGA GAC TGT GCA GAT GTA TAT CAA GCT GGT TTT AAT AAA AGT GGA
Arg Asp Cys Ala Asp Val Tyr Gln Ala Gly Phe Asn Lys Ser Gly>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      100     110     120     130
*   *   *   *   *   *   *   *
ATC TAC ACT ATT TAT ATT AAT AAT ATG CCA GAA CCC AAA AAG GTG
Ile Tyr Thr Ile Tyr Ile Asn Asn Met Pro Glu Pro Lys Lys Val>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      140     150     160     170     180
*   *   *   *   *   *   *   *
TTT TGC AAT ATG GAT GTC AAT GGG GGA GGT TGG ACT GTA ATA CAA
Phe Cys Asn Met Asp Val Asn Gly Gly Trp Thr Val Ile Gln>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      190     200     210     220
*   *   *   *   *   *   *   *
CAT CGT GAA GAT GGA AGT CTA GAT TTC CAA AGA GGC TGG AAG GAA
His Arg Glu Asp Gly Ser Leu Asp Phe Gln Arg Gly Trp Lys Glu>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      230     240     250     260     270
*   *   *   *   *   *   *   *
TAT AAA ATG GGT TTT GGA AAT CCC TCC GGT GAA TAT TGG CTG GGG
Tyr Lys Met Gly Phe Gly Asn Pro Ser Gly Glu Tyr Trp Leu Gly>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      280     290     300     310
*   *   *   *   *   *   *   *
AAT GAG TTT ATT TTT GCC ATT ACC AGT CAG AGG CAG TAC ATG CTA
Asn Glu Phe Ile Phe Ala Ile Thr Ser Gln Arg Gln Tyr Met Leu>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      320     330     340     350     360
*   *   *   *   *   *   *   *
AGA ATT GAG TTA ATG GAC TGG GAA GGG AAC CGA GCC TAT TCA CAG
Arg Ile Glu Leu Met Asp Trp Glu Gly Asn Arg Ala Tyr Ser Gln>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      370     380     390     400
*   *   *   *   *   *   *   *
TAT GAC AGA TTC CAC ATA GGA AAT GAA AAG CAA AAC TAT AGG TTG
Tyr Asp Arg Phe His Ile Gly Asn Glu Lys Gln Asn Tyr Arg Leu>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

      410     420     430     440     450
*   *   *   *   *   *   *   *
TAT TTA AAA GGT CAC ACT GGG ACA GCA GGA AAA CAG AGC AGC CTG
Tyr Leu Lys Gly His Thr Gly Thr Ala Gly Lys Gln Ser Ser Leu>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b>

```

2/42
Figure 1B

```

      460      470      480      490
      *      *      *      *
ATC TTA CAC GGT GCT GAT TTC AGC ACT AAA GAT GCT GAT AAT GAC
Ile Leu His Gly Ala Asp Phe Ser Thr Lys Asp Ala Asp Asn Asp>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      500      510      520      530      540
      *      *      *      *      *
AAC TGT ATG TGC AAA TGT GCC CTC ATG TTA ACA GGA GGA TGG TGG
Asn Cys Met Cys Lys Cys Ala Leu Met Leu Thr Gly Gly Trp Trp>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      550      560      570      580
      *      *      *      *
TTT GAT GCT TGT GGC CCC TCC AAT CTA AAT GGA ATG TTC TAT ACT
Phe Asp Ala Cys Gly Pro Ser Asn Leu Asn Gly Met Phe Tyr Thr>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      590      600      610      620      630
      *      *      *      *      *
GCG GGA CAA AAC CAT GGA AAA CTG AAT GGG ATA AAG TGG CAC TAC
Ala Gly Gln Asn His Gly Lys Leu Asn Gly Ile Lys Trp His Tyr>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      640      650      660      670
      *      *      *      *
TTC AAA GGG CCC AGT TAC TCC TTA CGT TCC ACA ACT ATG ATG ATT
Phe Lys Gly Pro Ser Tyr Ser Leu Arg Ser Thr Thr Met Met Ile>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      680      690      700      710      720
      *      *      *      *      *
CGA CCT TTA GAT TTT GGC CCC GCG CCT TTT AGA GAC TGT GCA GAT
Arg Pro Leu Asp Phe>
__ANG1 FIBRINO__>
      Gly Pro Ala Pro>
      __GPAP BRI__>
      Phe Arg Asp Cys Ala Asp>
      __ANG1 FIBRINOGEN-__>

      730      740      750      760
      *      *      *      *
GTA TAT CAA GCT GGT TTT AAT AAA AGT GGA ATC TAC ACT ATT TAT
Val Tyr Gln Ala Gly Phe Asn Lys Ser Gly Ile Tyr Thr Ile Tyr>
__d__d__d__ANG1 FIBRINOGEN-LIKE DOMAIN_d__d__d__d__>

      770      780      790      800      810
      *      *      *      *      *
ATT AAT AAT ATG CCA GAA CCC AAA AAG GTG TTT TGC AAT ATG GAT
Ile Asn Asn Met Pro Glu Pro Lys Lys Val Phe Cys Asn Met Asp>
__d__d__d__ANG1 FIBRINOGEN-LIKE DOMAIN_d__d__d__d__>

      820      830      840      850
      *      *      *      *
GTC AAT GGG GGA GGT TGG ACT GTA ATA CAA CAT CGT GAA GAT GGA
Val Asn Gly Gly Gly Trp Thr Val Ile Gln His Arg Glu Asp Gly>
__d__d__d__ANG1 FIBRINOGEN-LIKE DOMAIN_d__d__d__d__>

```


3/42
Figure 1C

```

      860      870      880      890      900
      *      *      *      *      *
AGT CTA GAT TTC CAA AGA GGC TGG AAG GAA TAT AAA ATG GGT TTT
Ser Leu Asp Phe Gln Arg Gly Trp Lys Glu Tyr Lys Met Gly Phe>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      910      920      930      940
      *      *      *      *      *
GGA AAT CCC TCC GGT GAA TAT TGG CTG GGG AAT GAG TTT ATT TTT
Gly Asn Pro Ser Gly Glu Tyr Trp Leu Gly Asn Glu Phe Ile Phe>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      950      960      970      980      990
      *      *      *      *      *
GCC ATT ACC AGT CAG AGG CAG TAC ATG CTA AGA ATT GAG TTA ATG
Ala Ile Thr Ser Gln Arg Gln Tyr Met Leu Arg Ile Glu Leu Met>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1000     1010     1020     1030
      *      *      *      *      *
GAC TGG GAA GGG AAC CGA GCC TAT TCA CAG TAT GAC AGA TTC CAC
Asp Trp Glu Gly Asn Arg Ala Tyr Ser Gln Tyr Asp Arg Phe His>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1040     1050     1060     1070     1080
      *      *      *      *      *
ATA GGA AAT GAA AAG CAA AAC TAT AGG TTG TAT TTA AAA GGT CAC
Ile Gly Asn Glu Lys Gln Asn Tyr Arg Leu Tyr Leu Lys Gly His>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1090     1100     1110     1120
      *      *      *      *      *
ACT GGG ACA GCA GGA AAA CAG AGC AGC CTG ATC TTA CAC GGT GCT
Thr Gly Thr Ala Gly Lys Gln Ser Ser Leu Ile Leu His Gly Ala>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1130     1140     1150     1160     1170
      *      *      *      *      *
GAT TTC AGC ACT AAA GAT GCT GAT AAT GAC AAC TGT ATG TGC AAA
Asp Phe Ser Thr Lys Asp Ala Asp Asn Asp Asn Cys Met Cys Lys>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1180     1190     1200     1210
      *      *      *      *      *
TGT GCC CTC ATG TTA ACA GGA GGA TGG TGG TTT GAT GCT TGT GGC
Cys Ala Leu Met Leu Thr Gly Gly Trp Trp Phe Asp Ala Cys Gly>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1220     1230     1240     1250     1260
      *      *      *      *      *
CCC TCC AAT CTA AAT GGA ATG TTC TAT ACT GCG GGA CAA AAC CAT
Pro Ser Asn Leu Asn Gly Met Phe Tyr Thr Ala Gly Gln Asn His>
_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

```

4/42
Figure 1D

```

      1270      1280      1290      1300
      *      *      *      *      *
GGA AAA CTG AAT GGG ATA AAG TGG CAC TAC TTC AAA GGG CCC AGT
Gly Lys Leu Asn Gly Ile Lys Trp His Tyr Phe Lys Gly Pro Ser>
_d_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1310      1320      1330      1340      1350
      *      *      *      *      *
TAC TCC TTA CGT TCC ACA ACT ATG ATG ATT CGA CCT TTA GAT TTT
Tyr Ser Leu Arg Ser Thr Thr Met Met Ile Arg Pro Leu Asp Phe>
_d_d_d_d_ANG1 FIBRINOGEN-LIKE DOMAIN_d_d_d_d_>

      1360      1370      1380      1390
      *      *      *      *      *
GGA CCG GGC GAG CCC AAA TCT TGT GAC AAA ACT CAC ACA TGC CCA
Gly Pro Gly>
_e_e_e_>
      Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro>
      _f_f_f_f_FC TAG (SPLIT)_f_f_f_f_>

      1400      1410      1420      1430      1440
      *      *      *      *      *
CCG TGC CCA GCA CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC CTC
Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu>
_f_f_f_f_f_FC TAG (SPLIT)_f_f_f_f_f_>

      1450      1460      1470      1480
      *      *      *      *      *
TTC CCC CCA AAA CCC AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT
Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro>
_f_f_f_f_f_FC TAG (SPLIT)_f_f_f_f_f_>

      1490      1500      1510      1520      1530
      *      *      *      *      *
GAG GTC ACA TGC GTG GTG GTG GAC GTG AGC CAC GAA GAC CCT GAG
Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu>
_f_f_f_f_f_FC TAG (SPLIT)_f_f_f_f_f_>

      1540      1550      1560      1570
      *      *      *      *      *
GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG GAG GTG CAT AAT GCC
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala>
_f_f_f_f_f_FC TAG (SPLIT)_f_f_f_f_f_>

      1580      1590      1600      1610      1620
      *      *      *      *      *
AAG ACA AAG CCG CGG GAG GAG CAG TAC AAC AGC ACG TAC CGT GTG
Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val>
_f_f_f_f_f_FC TAG (SPLIT)_f_f_f_f_f_>

      1630      1640      1650      1660
      *      *      *      *      *
GTC AGC GTC CTC ACC GTC CTG CAC CAG GAC TGG CTG AAT GGC AAG
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys>
_f_f_f_f_f_FC TAG (SPLIT)_f_f_f_f_f_>

```

5/42

Figure 1E

```

1670      1680      1690      1700      1710
*          *          *          *          *
GAG TAC AAG TGC AAG GTC TCC AAC AAA GCC CTC CCA GCC CCC ATC
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

      1720      1730      1740      1750
*          *          *          *          *
GAG AAA ACC ATC TCC AAA GCC AAA GGG CAG CCC CGA GAA CCA CAG
Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

1760      1770      1780      1790      1800
*          *          *          *          *
GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC AAG AAC CAG
Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

      1810      1820      1830      1840
*          *          *          *          *
GTC AGC CTG ACC TGC CTG GTC AAA GGC TTC TAT CCC AGC GAC ATC
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

1850      1860      1870      1880      1890
*          *          *          *          *
GCC GTG GAG TGG GAG AGC AAT GGG CAG CCG GAG AAC AAC TAC AAG
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

      1900      1910      1920      1930
*          *          *          *          *
ACC ACG CCT CCC GTG CTG GAC TCC GAC GGC TCC TTC TTC CTC TAC
Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

1940      1950      1960      1970      1980
*          *          *          *          *
AGC AAG CTC ACC GTG GAC AAG AGC AGG TGG CAG CAG GGG AAC GTC
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

      1990      2000      2010      2020
*          *          *          *          *
TTC TCA TGC TCC GTG ATG CAT GAG GCT CTG CAC AAC CAC TAC ACG
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

2030      2040      2050
*          *          *          *          *
CAG AAG AGC CTC TCC CTG TCT CCG GGT AAA TGA
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys ***>
__f__f__f__f__f__FC TAG (SPLIT)__f__f__f__f__f__>

```

Figure 2A

Figure 1A

10 20 30 40

ATG TCT GCA CTT CTG ATC CTA GCT CTT GTT GGA GCT GCA GTT GCT
Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val Ala>
_a_a_a_a_TRYPSIN SIGNAL SEQUENCE_a_a_a_a_a>

50 60 70 80 90

AGA GAC TGT GCT GAA GTA TTC AAA TCA GGA CAC ACC ACA AAT GGC
Arg Asp Cys Ala Glu Val Phe Lys Ser Gly His Thr Thr Asn Gly>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

100 110 120 130

ATC TAC ACG TTA ACA TTC CCT AAT TCT ACA GAA GAG ATC AAG GCC
Ile Tyr Thr Leu Thr Phe Pro Asn Ser Thr Glu Glu Ile Lys Ala>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

140 150 160 170 180

TAC TGT GAC ATG GAA GCT GGA GGA GGC TGG ACA ATT ATT CAG
Tyr Cys Asp Met Glu Ala Gly Gly Gly Trp Thr Thr Ile Ile Gln>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

190 200 210 220

CGA CGT GAG GAT GGC AGC GTT GAT TTT CAG AGG ACT TGG AAA GAA
Arg Arg Glu Asp Gly Ser Val Asp Phe Gln Arg Thr Trp Lys Glu>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

230 240 250 260 270

TAT AAA GTG GGA TTT GGT AAC CCT TCA GGA GAA TAT TGG CTG GGA
Tyr Lys Val Gly Phe Gly Asn Pro Ser Gly Glu Tyr Trp Leu Gly>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

280 290 300 310

AAT GAG TTT GTT TCG CAA CTG ACT AAT CAG CAA CGC TAT GTG CTT
Asn Glu Phe Val Ser Gln Leu Thr Asn Gln Gln Arg Tyr Val Leu>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

320 330 340 350 360

AAA ATA CAC CTT AAA GAC TGG GAA GGG AAT GAG GCT TAC TCA TTG
Lys Ile His Leu Lys Asp Trp Glu Gly Asn Glu Ala Tyr Ser Leu>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

370 380 390 400

TAT GAA CAT TTC TAT CTC TCA AGT GAA GAA CTC AAT TAT AGG ATT
Tyr Glu His Phe Tyr Leu Ser Ser Glu Glu Leu Asn Tyr Arg Ile>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

410 420 430 440 450

CAC CTT AAA GGA CTT ACA GGG ACA GCC GGC AAA ATA AGC AGC ATC
His Leu Lys Gly Leu Thr Gly Thr Ala Gly Lys Ile Ser Ser Ile>
_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN #1_b_b_b_b>

7/42
Figure 2B

```

      460      470      480      490
      *      *      *      *
AGC CAA CCA GGA AAT GAT TTT AGC ACA AAG GAT GGA GAC AAC GAC
Ser Gln Pro Gly Asn Asp Phe Ser Thr Lys Asp Gly Asp Asn Asp>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN #1__b__b__b__>

      500      510      520      530      540
      *      *      *      *      *
AAA TGT ATT TGC AAA TGT TCA CAA ATG CTA ACA GGA GGC TGG TGG
Lys Cys Ile Cys Lys Cys Ser Gln Met Leu Thr Gly Gly Trp Trp>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN #1__b__b__b__>

      550      560      570      580
      *      *      *      *
TTT GAT GCA TGT GGT CCT TCC AAC TTG AAC GGA ATG TAC TAT CCA
Phe Asp Ala Cys Gly Pro Ser Asn Leu Asn Gly Met Tyr Tyr Pro>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN #1__b__b__b__>

      590      600      610      620      630
      *      *      *      *      *
CAG AGG CAG AAC ACA AAT AAG TTC AAC GGC ATT AAA TGG TAC TAC
Gln Arg Gln Asn Thr Asn Lys Phe Asn Gly Ile Lys Trp Tyr Tyr>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN #1__b__b__b__>

      640      650      660      670
      *      *      *      *
TGG AAA GGC TCA GGC TAT TCG CTC AAG GCC ACA ACC ATG ATG ATC
Trp Lys Gly Ser Gly Tyr Ser Leu Lys Ala Thr Thr Met Met Ile>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN #1__b__b__b__>

      680      690      700      710      720
      *      *      *      *      *
CGA CCA GCA GAT TTC GGG GGC CCC GCG CCT TTC AGA GAC TGT GCT
Arg Pro Ala Asp Phe>
__ANG2 FIBRINO__>
      Gly Gly Pro Ala Pro>
      __GGPAP BRIDGE__>
      Phe Arg Asp Cys Ala>
      __ANG2 FIBRINO__>

      730      740      750      760
      *      *      *      *
GAA GTA TTC AAA TCA GGA CAC ACC ACA AAT GGC ATC TAC ACG TTA
Glu Val Phe Lys Ser Gly His Thr Thr Asn Gly Ile Tyr Thr Leu>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      770      780      790      800      810
      *      *      *      *      *
ACA TTC CCT AAT TCT ACA GAA GAG ATC AAG GCC TAC TGT GAC ATG
Thr Phe Pro Asn Ser Thr Glu Glu Ile Lys Ala Tyr Cys Asp Met>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      820      830      840      850
      *      *      *      *
GAA GCT GGA GGA GGC GGG TGG ACA ATT ATT CAG CGA CGT GAG GAT
Glu Ala Gly Gly Gly Trp Thr Ile Ile Gln Arg Arg Glu Asp>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

```

8/42
Figure 2C

```

      860      870      880      890      900
      *      *      *      *      *
GGC AGC GTT GAT TTT CAG AGG ACT TGG AAA GAA TAT AAA GTG GGA
Gly Ser Val Asp Phe Gln Arg Thr Trp Lys Glu Tyr Lys Val Gly>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      910      920      930      940
      *      *      *      *
TTT GGT AAC CCT TCA GGA GAA TAT TGG CTG GGA AAT GAG TTT GTT
Phe Gly Asn Pro Ser Gly Glu Tyr Trp Leu Gly Asn Glu Phe Val>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      950      960      970      980      990
      *      *      *      *      *
TCG CAA CTG ACT AAT CAG CAA CGC TAT GTG CTT AAA ATA CAC CTT
Ser Gln Leu Thr Asn Gln Gln Arg Tyr Val Leu Lys Ile His Leu>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      1000     1010     1020     1030
      *      *      *      *
AAA GAC TGG GAA GGG AAT GAG GCT TAC TCA TTG TAT GAA CAT TTC
Lys Asp Trp Glu Gly Asn Glu Ala Tyr Ser Leu Tyr Glu His Phe>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      1040     1050     1060     1070     1080
      *      *      *      *      *
TAT CTC TCA AGT GAA GAA CTC AAT TAT AGG ATT CAC CTT AAA GGA
Tyr Leu Ser Ser Glu Glu Leu Asn Tyr Arg Ile His Leu Lys Gly>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      1090     1100     1110     1120
      *      *      *      *
CTT ACA GGG ACA GCC GGC AAA ATA AGC AGC ATC AGC CAA CCA GGA
Leu Thr Gly Thr Ala Gly Lys Ile Ser Ser Ile Ser Gln Pro Gly>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      1130     1140     1150     1160     1170
      *      *      *      *      *
AAT GAT TTT AGC ACA AAG GAT GGA GAC AAC GAC AAA TGT ATT TGC
Asn Asp Phe Ser Thr Lys Asp Gly Asp Asn Asp Lys Cys Ile Cys>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      1180     1190     1200     1210
      *      *      *      *
AAA TGT TCA CAA ATG CTA ACA GGA GGC TGG TGG TTT GAT GCA TGT
Lys Cys Ser Gln Met Leu Thr Gly Gly Trp Trp Phe Asp Ala Cys>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

      1220     1230     1240     1250     1260
      *      *      *      *      *
GGT CCT TCC AAC TTG AAC GGA ATG TAC TAT CCA CAG AGG CAG AAC
Gly Pro Ser Asn Leu Asn Gly Met Tyr Tyr Pro Gln Arg Gln Asn>
__d__d__d__ANG2 FIBRINOGEN-LIKE DOMAIN#2__d__d__d__>

```

9/42
Figure 2D

```

      1270      1280      1290      1300
      *      *      *      *
ACA AAT AAG TTC AAC GGC ATT AAA TGG TAC TAC TGG AAA GGC TCA
Thr Asn Lys Phe Asn Gly Ile Lys Trp Tyr Tyr Trp Lys Gly Ser>
_d_d_d_ANG2 FIBRINOGEN-LIKE DOMAIN#2_d_d_d_>

      1310      1320      1330      1340      1350
      *      *      *      *      *
GGC TAT TCG CTC AAG GCC ACA ACC ATG ATG ATC CGA CCA GCA GAT
Gly Tyr Ser Leu Lys Ala Thr Thr Met Met Ile Arg Pro Ala Asp>
_d_d_d_ANG2 FIBRINOGEN-LIKE DOMAIN#2_d_d_d_>

      1360      1370      1380      1390
      *      *      *      *
TTC GGA CCG GGC GAG CCC AAA TCT TGT GAC AAA ACT CAC ACA TGC
Phe>
____>
      Gly Pro Gly>
      _e_e_>
      Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys>
      _f_f_f_f_FC TAG_f_f_f_f_>

      1400      1410      1420      1430      1440
      *      *      *      *      *
CCA CCG TGC CCA GCA CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC
Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_>

      1450      1460      1470      1480
      *      *      *      *
CTC TTC CCC CCA AAA CCC AAG GAC ACC CTC ATG ATC TCC CGG ACC
Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_>

      1490      1500      1510      1520      1530
      *      *      *      *      *
CCT GAG GTC ACA TGC GTG GTG GTG GAC GTG AGC CAC GAA GAC CCT
Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_>

      1540      1550      1560      1570
      *      *      *      *
GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG GAG GTG CAT AAT
Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_>

      1580      1590      1600      1610      1620
      *      *      *      *      *
GCC AAG ACA AAG CCG CGG GAG GAG CAG TAC AAC AGC ACG TAC CGT
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_>

      1630      1640      1650      1660
      *      *      *      *
GTG GTC AGC GTC CTC ACC GTC CTG CAC CAG GAC TGG CTG AAT GGC
Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_>

```

10/42
Figure 2E

1670 1680 1690 1700 1710
AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA GCC CTC CCA GCC CCC
Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

1720 1730 1740 1750
ATC GAG AAA ACC ATC TCC AAA GCC AAA GGG CAG CCC CGA GAA CCA
Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

1760 1770 1780 1790 1800
CAG GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC AAG AAC
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

1810 1820 1830 1840
CAG GTC AGC CTG ACC TGC CTG GTC AAA GGC TTC TAT CCC AGC GAC
Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

1850 1860 1870 1880 1890
ATC GCC GTG GAG TGG GAG AGC AAT GGG CAG CCG GAG AAC AAC TAC
Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

1900 1910 1920 1930
AAG ACC ACG CCT CCC GTG CTG GAC TCC GAC GGC TCC TTC TTC CTC
Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

1940 1950 1960 1970 1980
TAC AGC AAG CTC ACC GTG GAC AAG AGC AGG TGG CAG CAG GGG AAC
Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

1990 2000 2010 2020
GTC TTC TCA TGC TCC GTG ATG CAT GAG GCT CTG CAC AAC CAC TAC
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>

2030 2040 2050 2060
ACG CAG AAG AGC CTC TCC CTG TCT CCG GGT AAA TGA
Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys ***>
_f_f_f_f_f_f_FC TAG_f_f_f_f_f_f_f_f_>


```

      10      20      30      40
      *      *      *      *
ATG TCT GCA CTT CTG ATC CTA GCT CTT GTT GGA GCT GCA GTT GCT
Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val-Ala>
_a_a_a_a_TRYPSIN SIGNAL SEQUENCE_a_a_a_a>

      50      60      70      80      90
      *      *      *      *      *
AGA GAC TGT GCA GAT GTA TAT CAA GCT GGT TTT AAT AAA AGT GGA
Arg Asp Cys Ala Asp Val Tyr Gln Ala Gly Phe Asn Lys Ser Gly>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

      100     110     120     130
      *      *      *      *      *
ATC TAC ACT ATT TAT ATT AAT AAT ATG CCA GAA CCC AAA AAG GTG
Ile Tyr Thr Ile Tyr Ile Asn Asn Met Pro Glu Pro Lys Lys Val>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

      140     150     160     170     180
      *      *      *      *      *
TTT TGC AAT ATG GAT GTC AAT GGG GGA GGT TGG ACT GTA ATA CAA
Phe Cys Asn Met Asp Val Asn Gly Gly Gly Trp Thr Val Ile Gln>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

      190     200     210     220
      *      *      *      *      *
CAT CGT GAA GAT GGA AGT CTA GAT TTC CAA AGA GGC TGG AAG GAA
His Arg Glu Asp Gly Ser Leu Asp Phe Gln Arg Gly Trp Lys Glu>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

      230     240     250     260     270
      *      *      *      *      *
TAT AAA ATG GGT TTT GGA AAT CCC TCC GGT GAA TAT TGG CTG GGG
Tyr Lys Met Gly Phe Gly Asn Pro Ser Gly Glu Tyr Trp Leu Gly>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

      280     290     300     310
      *      *      *      *      *
AAT GAG TTT ATT TTT GCC ATT ACC AGT CAG AGG CAG TAC ATG CTA
Asn Glu Phe Ile Phe Ala Ile Thr Ser Gln Arg Gln Tyr Met Leu>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

      320     330     340     350     360
      *      *      *      *      *
AGA ATT GAG TTA ATG GAC TGG GAA GGG AAC CGA GCC TAT TCA CAG
Arg Ile Glu Leu Met Asp Trp Glu Gly Asn Arg Ala Tyr Ser Gln>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

      370     380     390     400
      *      *      *      *      *
TAT GAC AGA TTC CAC ATA GGA AAT GAA AAG CAA AAC TAT AGG TTG
Tyr Asp Arg Phe His Ile Gly Asn Glu Lys Gln Asn Tyr Arg Leu>
_b_b_b_b_ANG1 FIBRINOGEN-LIKE DOMAIN_b_b_b_b>

```

12/42
Figure 3B

```

      410      420      430      440      450
      *      *      *      *      *
TAT TTA AAA GGT CAC ACT GGG ACA GCA GGA AAA CAG AGC AGC CTG
Tyr Leu Lys Gly His Thr Gly Thr Ala Gly Lys Gln Ser Ser Leu>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      460      470      480      490
      *      *      *      *      *
ATC TTA CAC GGT GCT GAT TTC AGC ACT AAA GAT GCT GAT AAT GAC
Ile Leu His Gly Ala Asp Phe Ser Thr Lys Asp Ala Asp Asn Asp>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      500      510      520      530      540
      *      *      *      *      *
AAC TGT ATG TGC AAA TGT GCC CTC ATG TTA ACA GGA GGA TGG TGG
Asn Cys Met Cys Lys Cys Ala Leu Met Leu Thr Gly Gly Trp Trp>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      550      560      570      580
      *      *      *      *      *
TTT GAT GCT TGT GGC CCC TCC AAT CTA AAT GGA ATG TTC TAT ACT
Phe Asp Ala Cys Gly Pro Ser Asn Leu Asn Gly Met Phe Tyr Thr>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      590      600      610      620      630
      *      *      *      *      *
GCG GGA CAA AAC CAT GGA AAA CTG AAT GGG ATA AAG TGG CAC TAC
Ala Gly Gln Asn His Gly Lys Leu Asn Gly Ile Lys Trp His Tyr>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      640      650      660      670
      *      *      *      *      *
TTC AAA GGG CCA AGT TAC TCC TTA CGT TCC ACA ACT ATG ATG ATT
Phe Lys Gly Pro Ser Tyr Ser Leu Arg Ser Thr Thr Met Met Ile>
__b__b__b__ANG1 FIBRINOGEN-LIKE DOMAIN_b__b__b__b__>

      680      690      700      710      720
      *      *      *      *      *
CGA CCT TTA GAT TTT GGC CCG GGC GAG CCC AAA TCT TGT GAC AAA
Arg Pro Leu Asp Phe>
__ANG1 FIBRINO__>
      Gly Pro Gly>
      __c__c__>
      Glu Pro Lys Ser Cys Asp Lys>
      __d__d__FC TAG__d__d__>

      730      740      750      760
      *      *      *      *      *
ACT CAC ACA TGC CCA CCG TGC CCA GCA CCT GAA CTC CTG GGG GGA
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly>
__d__d__d__d__d__d__FC TAG__d__d__d__d__d__d__>

      770      780      790      800      810
      *      *      *      *      *
CCG TCA GTC TTC CTC TTC CCC CCA AAA CCC AAG GAC ACC CTC ATG
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met>
__d__d__d__d__d__d__FC TAG__d__d__d__d__d__d__>

```

13/42
Figure 3C

```

      820      830      840      850
      *      *      *      *
ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG GTG GAC GTG AGC
Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

      860      870      880      890      900
      *      *      *      *      *
CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG
His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

      910      920      930      940
      *      *      *      *      *
GAG GTG CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG TAC AAC
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

      950      960      970      980      990
      *      *      *      *      *
AGC ACG TAC CGT GTG GTC AGC GTC CTC ACC GTC CTG CAC CAG GAC
Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

     1000     1010     1020     1030
      *      *      *      *      *
TGG CTG AAT GGC AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA GCC
Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

     1040     1050     1060     1070     1080
      *      *      *      *      *
CTC CCA GCC CCC ATC GAG AAA ACC ATC TCC AAA GCC AAA GGG CAG
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

     1090     1100     1110     1120
      *      *      *      *      *
CCC CGA GAA CCA CAG GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

     1130     1140     1150     1160     1170
      *      *      *      *      *
CTG ACC AAG AAC CAG GTC AGC CTG ACC TGC CTG GTC AAA GGC TTC
Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

     1180     1190     1200     1210
      *      *      *      *      *
TAT CCC AGC GAC ATC GCC GTG GAG TGG GAG AGC AAT GGG CAG CCG
Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

     1220     1230     1240     1250     1260
      *      *      *      *      *
GAG AAC AAC TAC AAG ACC ACG CCT CCC GTG CTG GAC TCC GAC GGC
Glu Asn Asn Tyr Lys Thr Pro Pro Val Leu Asp Ser Asp Gly>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d>

```

14/42
Figure 3D

```

      1270      1280      1290      1300
      *      *      *      *
TCC TTC TTC CTC TAC AGC AAG CTC ACC GTG GAC AAG AGC AGG TGG
Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_>

      1310      1320      1330      1340      1350
      *      *      *      *      *
CAG CAG GGG AAC GTC TTC TCA TGC TCC GTG ATG CAT GAG GCT CTG
Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_>

      1360      1370      1380      1390
      *      *      *      *
CAC AAC CAC TAC ACG CAG AAG AGC CTC TCC CTG TCT CCG GGT AAA
His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_>

      1400      1410      1420      1430      1440
      *      *      *      *      *
GGC GGT GGC GGT TCT GGC GCG CCT TTT AGA GAC TGT GCA GAT GTA
Gly Gly Gly Gly Ser Gly Ala Pro>
_G4S LINKER/ASC BRIDGE (N____>
                                Phe Arg Asp Cys Ala Asp Val>
                                _ANG1 FIBRINOGEN-LIKE____>

      1450      1460      1470      1480
      *      *      *      *
TAT CAA GCT GGT TTT AAT AAA AGT GGA ATC TAC ACT ATT TAT ATT
Tyr Gln Ala Gly Phe Asn Lys Ser Gly Ile Tyr Thr Ile Tyr Ile>
_f_f_f_f_ANG1 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1490      1500      1510      1520      1530
      *      *      *      *      *
AAT AAT ATG CCA GAA CCC AAA AAG GTG TTT TGC AAT ATG GAT GTC
Asn Asn Met Pro Glu Pro Lys Lys Val Phe Cys Asn Met Asp Val>
_f_f_f_f_ANG1 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1540      1550      1560      1570
      *      *      *      *
AAT GGG GGA GGT TGG ACT GTA ATA CAA CAT CGT GAA GAT GGA AGT
Asn Gly Gly Gly Trp Thr Val Ile Gln His Arg Glu Asp Gly Ser>
_f_f_f_f_ANG1 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1580      1590      1600      1610      1620
      *      *      *      *      *
CTA GAT TTC CAA AGA GGC TGG AAG GAA TAT AAA ATG GGT TTT GGA
Leu Asp Phe Gln Arg Gly Trp Lys Glu Tyr Lys Met Gly Phe Gly>
_f_f_f_f_ANG1 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1630      1640      1650      1660
      *      *      *      *
AAT CCC TCC GGT GAA TAT TGG CTG GGG AAT GAG TTT ATT TTT GCC
Asn Pro Ser Gly Glu Tyr Trp Leu Gly Asn Glu Phe Ile Phe Ala>
_f_f_f_f_ANG1 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

```

15/42
Figure 3E

```

1670      1680      1690      1700      1710
*          *          *          *          *
ATT ACC AGT CAG AGG CAG TAC ATG CTA AGA ATT GAG TTA ATG GAC
Ile Thr Ser Gln Arg Gln Tyr Met Leu Arg Ile Glu Leu Met Asp>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

      1720      1730      1740      1750
*          *          *          *          *
TGG GAA GGG AAC CGA GCC TAT TCA CAG TAT GAC AGA TTC CAC ATA
Trp Glu Gly Asn Arg Ala Tyr Ser Gln Tyr Asp Arg Phe His Ile>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

1760      1770      1780      1790      1800
*          *          *          *          *
GGA AAT GAA AAG CAA AAC TAT AGG TTG TAT TTA AAA GGT CAC ACT
Gly Asn Glu Lys Gln Asn Tyr Arg Leu Tyr Leu Lys Gly His Thr>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

      1810      1820      1830      1840
*          *          *          *          *
GGG ACA GCA GGA AAA CAG AGC AGC CTG ATC TTA CAC GGT GCT GAT
Gly Thr Ala Gly Lys Gln Ser Ser Leu Ile Leu His Gly Ala Asp>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

1850      1860      1870      1880      1890
*          *          *          *          *
TTC AGC ACT AAA GAT GCT GAT AAT GAC AAC TGT ATG TGC AAA TGT
Phe Ser Thr Lys Asp Ala Asp Asn Asp Asn Cys Met Cys Lys Cys>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

      1900      1910      1920      1930
*          *          *          *          *
GCC CTC ATG TTA ACA GGA GGA TGG TGG TTT GAT GCT TGT GGC CCC
Ala Leu Met Leu Thr Gly Gly Trp Trp Phe Asp Ala Cys Gly Pro>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

1940      1950      1960      1970      1980
*          *          *          *          *
TCC AAT CTA AAT GGA ATG TTC TAT ACT GCG GGA CAA AAC CAT GGA
Ser Asn Leu Asn Gly Met Phe Tyr Thr Ala Gly Gln Asn His Gly>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

      1990      2000      2010      2020
*          *          *          *          *
AAA CTG AAT GGG ATA AAG TGG CAC TAC TTC AAA GGG CCA AGT TAC
Lys Leu Asn Gly Ile Lys Trp His Tyr Phe Lys Gly Pro Ser Tyr>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

2030      2040      2050      2060
*          *          *          *          *
TCC TTA CGT TCC ACA ACT ATG ATG ATT CGA CCT TTA GAT TTT
Ser Leu Arg Ser Thr Thr Met Met Ile Arg Pro Leu Asp Phe>
__f__f__f__ANG1 FIBRINOGEN-LIKE DOMAIN__f__f__f__f__>

```

16/42

Figure 4A

```

      10      20      30      40
      *      *      *      *
ATG TCT GCA CTT CTG ATC CTA GCT CTT GTT GGA GCT GCA GTT GCT
Met Ser Ala Leu Leu Ile Leu Ala Leu Val Gly Ala Ala Val Ala>
_a_a_a_a_a_TRYPSIN SIGNAL SEQUENCE_a_a_a_a_a_>

      50      60      70      80      90
      *      *      *      *      *
AGA GAC TGT GCT GAA GTA TTC AAA TCA GGA CAC ACC ACA AAT GGC
Arg Asp Cys Ala Glu Val Phe Lys Ser Gly His Thr Thr Asn Gly>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

      100     110     120     130
      *      *      *      *      *
ATC TAC ACG TTA ACA TTC CCT AAT TCT ACA GAA GAG ATC AAG GCC
Ile Tyr Thr Leu Thr Phe Pro Asn Ser Thr Glu Glu Ile Lys Ala>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

      140     150     160     170     180
      *      *      *      *      *
TAC TGT GAC ATG GAA GCT GGA GGA GGC GGG TGG ACA ATT ATT CAG
Tyr Cys Asp Met Glu Ala Gly Gly Gly Gly Trp Thr Ile Ile Gln>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

      190     200     210     220
      *      *      *      *      *
CGA CGT GAG GAT GGC AGC GTT GAT TTT CAG AGG ACT TGG AAA GAA
Arg Arg Glu Asp Gly Ser Val Asp Phe Gln Arg Thr Trp Lys Glu>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

      230     240     250     260     270
      *      *      *      *      *
TAT AAA GTG GGA TTT GGT AAC CCT TCA GGA GAA TAT TGG CTG GGA
Tyr Lys Val Gly Phe Gly Asn Pro Ser Gly Glu Tyr Trp Leu Gly>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

      280     290     300     310
      *      *      *      *      *
AAT GAG TTT GTT TCG CAA CTG ACT AAT CAG CAA CGC TAT GTG CTT
Asn Glu Phe Val Ser Gln Leu Thr Asn Gln Gln Arg Tyr Val Leu>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

      320     330     340     350     360
      *      *      *      *      *
AAA ATA CAC CTT AAA GAC TGG GAA GGG AAT GAG GCT TAC TCA TTG
Lys Ile His Leu Lys Asp Trp Glu Gly Asn Glu Ala Tyr Ser Leu>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

      370     380     390     400
      *      *      *      *      *
TAT GAA CAT TTC TAT CTC TCA AGT GAA GAA CTC AAT TAT AGG ATT
Tyr Glu His Phe Tyr Leu Ser Ser Glu Glu Leu Asn Tyr Arg Ile>
_b_b_b_b_ANG2 FIBRINOGEN-LIKE DOMAIN_b_b_b_b_b_>

```

17/42
Figure 4B

```

      410      420      430      440      450
      *      *      *      *      *
CAC CTT AAA GGA CTT ACA GGG ACA GCC GGC AAA ATA AGC AGC ATC
His Leu Lys Gly Leu Thr Gly Thr Ala Gly Lys Ile Ser Ser Ile>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN__b__b__b__b__>

      460      470      480      490
      *      *      *      *      *
AGC CAA CCA GGA AAT GAT TTT AGC ACA AAG GAT GGA GAC AAC GAC
Ser Gln Pro Gly Asn Asp Phe Ser Thr Lys Asp Gly Asp Asn Asp>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN__b__b__b__b__>

      500      510      520      530      540
      *      *      *      *      *
AAA TGT ATT TGC AAA TGT TCA CAA ATG CTA ACA GGA GGC TGG TGG
Lys Cys Ile Cys Lys Cys Ser Gln Met Leu Thr Gly Gly Trp Trp>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN__b__b__b__b__>

      550      560      570      580
      *      *      *      *      *
TTT GAT GCA TGT GGT CCT TCC AAC TTG AAC GGA ATG TAC TAT CCA
Phe Asp Ala Cys Gly Pro Ser Asn Leu Asn Gly Met Tyr Tyr Pro>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN__b__b__b__b__>

      590      600      610      620      630
      *      *      *      *      *
CAG AGG CAG AAC ACA AAT AAG TTC AAC GGC ATT AAA TGG TAC TAC
Gln Arg Gln Asn Thr Asn Lys Phe Asn Gly Ile Lys Trp Tyr Tyr>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN__b__b__b__b__>

      640      650      660      670
      *      *      *      *      *
TGG AAA GGC TCA GGC TAT TCG CTC AAG GCC ACA ACC ATG ATG ATC
Trp Lys Gly Ser Gly Tyr Ser Leu Lys Ala Thr Thr Met Met Ile>
__b__b__b__ANG2 FIBRINOGEN-LIKE DOMAIN__b__b__b__b__>

      680      690      700      710      720
      *      *      *      *      *
CGA CCA GCA GAT TTC GGG GGC CCG GGC GAG CCC AAA TCT TGT GAC
Arg Pro Ala Asp Phe>
__ANG2 FIBRINO__>
      Gly Gly Pro Gly>
      __GGPG BRI__>
      Glu Pro Lys Ser Cys Asp>
      __d__FC TAG__d__d__>

      730      740      750      760
      *      *      *      *      *
AAA ACT CAC ACA TGC CCA CCG TGC CCA GCA CCT GAA CTC CTG GGG
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly>
__d__d__d__d__d__d__FC TAG__d__d__d__d__d__d__>

      770      780      790      800      810
      *      *      *      *      *
GGA CCG TCA GTC TTC CTC TTC CCC CCA AAA CCC AAG GAC ACC CTC
Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu>
__d__d__d__d__d__d__FC TAG__d__d__d__d__d__d__>

```

18/42
Figure 4C

```

      820      830      840      850
      *      *      *      *
ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG GTG GAC GTG
Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      860      870      880      890      900
      *      *      *      *      *
AGC CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      910      920      930      940
      *      *      *      *      *
GTG GAG GTG CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG TAC
Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      950      960      970      980      990
      *      *      *      *      *
AAC AGC ACG TAC CGT GTG GTC AGC GTC CTC ACC GTC CTG CAC CAG
Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      1000     1010     1020     1030
      *      *      *      *      *
GAC TGG CTG AAT GGC AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      1040     1050     1060     1070     1080
      *      *      *      *      *
GCC CTC CCA GCC CCC ATC GAG AAA ACC ATC TCC AAA GCC AAA GGC
Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      1090     1100     1110     1120
      *      *      *      *      *
CAG CCC CGA GAA CCA CAG GTG TAC ACC CTG CCC CCA TCC CGG GAT
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Asp>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      1130     1140     1150     1160     1170
      *      *      *      *      *
GAG CTG ACC AAG AAC CAG GTC AGC CTG ACC TGC CTG GTC AAA GGC
Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      1180     1190     1200     1210
      *      *      *      *      *
TTC TAT CCC AGC GAC ATC GCC GTG GAG TGG GAG AGC AAT GGG CAG
Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

      1220     1230     1240     1250     1260
      *      *      *      *      *
CCG GAG AAC AAC TAC AAG ACC ACG CCT CCC GTG CTG GAC TCC GAC
Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp>
_d_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_d_>

```


19/42
Figure 4D

```

      1270      1280      1290      1300
      *          *          *          *
GGC TCC TTC TTC CTC TAC AGC AAG CTC ACC GTG GAC AAG AGC AGG
Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg>
_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_>

      1310      1320      1330      1340      1350
      *          *          *          *          *
TGG CAG CAG GGG AAC GTC TTC TCA TGC TCC GTG ATG CAT GAG GCT
Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala>
_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_>

      1360      1370      1380      1390
      *          *          *          *          *
CTG CAC AAC CAC TAC ACG CAG AAG AGC CTC TCC CTG TCT CCG GGT
Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly>
_d_d_d_d_d_d_d_FC TAG_d_d_d_d_d_d_d_d_d_>

      1400      1410      1420      1430      1440
      *          *          *          *          *
AAA GGC GGT GGC GGT TCT GGC GCG CCT AGA GAC TGT GCT GAA GTA
Lys>
____>
      Gly Gly Gly Gly Ser Gly Ala Pro>
      _e_GGGGSGAP BRIDGE_e____>
                                     Arg Asp Cys Ala Glu Val>
                                     _ANG2 FIBRINOGEN-____>

      1450      1460      1470      1480
      *          *          *          *          *
TTC AAA TCA GGA CAC ACC ACA AAT GGC ATC TAC ACG TTA ACA TTC
Phe Lys Ser Gly His Thr Thr Asn Gly Ile Tyr Thr Leu Thr Phe>
_f_f_f_f_ANG2 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1490      1500      1510      1520      1530
      *          *          *          *          *
CCT AAT TCT ACA GAA GAG ATC AAG GCC TAC TGT GAC ATG GAA GCT
Pro Asn Ser Thr Glu Glu Ile Lys Ala Tyr Cys Asp Met Glu Ala>
_f_f_f_f_ANG2 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1540      1550      1560      1570
      *          *          *          *          *
GGA GGA GGC GGG TGG ACA ATT ATT CAG CGA CGT GAG GAT GGC AGC
Gly Gly Gly Gly Trp Thr Ile Ile Gln Arg Arg Glu Asp Gly Ser>
_f_f_f_f_ANG2 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1580      1590      1600      1610      1620
      *          *          *          *          *
GTT GAT TTT CAG AGG ACT TGG AAA GAA TAT AAA GTG GGA TTT GGT
Val Asp Phe Gln Arg Thr Trp Lys Glu Tyr Lys Val Gly Phe Gly>
_f_f_f_f_ANG2 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

      1630      1640      1650      1660
      *          *          *          *          *
AAC CCT TCA GGA GAA TAT TGG CTG GGA AAT GAG TTT GTT TCG CAA
Asn Pro Ser Gly Glu Tyr Trp Leu Gly Asn Glu Phe Val Ser Gln>
_f_f_f_f_ANG2 FIBRINOGEN-LIKE DOMAIN_f_f_f_f_f_>

```

20/42
Figure 4E

```

1670      1680      1690      1700      1710
*          *          *          *          *
CTG ACT AAT CAG CAA CGC TAT GTG CTT AAA ATA CAC CTT AAA GAC
Leu Thr Asn Gln Gln Arg Tyr Val Leu Lys Ile His Leu Lys Asp>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

      1720      1730      1740      1750
*          *          *          *          *
TGG GAA GGG AAT GAG GCT TAC TCA TTG TAT GAA CAT TTC TAT CTC
Trp Glu Gly Asn Glu Ala Tyr Ser Leu Tyr Glu His Phe Tyr Leu>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

1760      1770      1780      1790      1800
*          *          *          *          *
TCA AGT GAA GAA CTC AAT TAT AGG ATT CAC CTT AAA GGA CTT ACA
Ser Ser Glu Glu Leu Asn Tyr Arg Ile His Leu Lys Gly Leu Thr>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

      1810      1820      1830      1840
*          *          *          *          *
GGG ACA GCC GGC AAA ATA AGC AGC ATC AGC CAA CCA GGA AAT GAT
Gly Thr Ala Gly Lys Ile Ser Ser Ile Ser Gln Pro Gly Asn Asp>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

1850      1860      1870      1880      1890
*          *          *          *          *
TTT AGC ACA AAG GAT GGA GAC AAC GAC AAA TGT ATT TGC AAA TGT
Phe Ser Thr Lys Asp Gly Asp Asn Asp Lys Cys Ile Cys Lys Cys>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

      1900      1910      1920      1930
*          *          *          *          *
TCA CAA ATG CTA ACA GGA GGC TGG TGG TTT GAT GCA TGT GGT CCT
Ser Gln Met Leu Thr Gly Gly Trp Trp Phe Asp Ala Cys Gly Pro>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

1940      1950      1960      1970      1980
*          *          *          *          *
TCC AAC TTG AAC GGA ATG TAC TAT CCA CAG AGG CAG AAC ACA AAT
Ser Asn Leu Asn Gly Met Tyr Tyr Pro Gln Arg Gln Asn Thr Asn>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>


      1990      2000      2010      2020
*          *          *          *          *
AAG TTC AAC GGC ATT AAA TGG TAC TAC TGG AAA GGC TCA GGC TAT
Lys Phe Asn Gly Ile Lys Trp Tyr Tyr Trp Lys Gly Ser Gly Tyr>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN_f__f__f__f__>

2030      2040      2050      2060      2070
*          *          *          *          *
TCG CTC AAG GCC ACA ACC ATG ATG ATC CGA CCA GCA GAT TTC TGA
Ser Leu Lys Ala Thr Thr Met Met Ile Arg Pro Ala Asp Phe>
__f__f__f__ANG2 FIBRINOGEN-LIKE DOMAIN__f__f__f__f__>

```

21/42


Figure 5



Ang1-FD-Fc-FD


22/42

Figure 6
Molar Mass vs. Volume




23/42

Figure 7




24/42

Figure 8




25/42

Figure 9




26/42

Figure 10




27/42

Figure 11




28/42

Figure 12



29/42

Figure 13



30/42

Figure 14A

```

      10      20      30      40
      *      *      *      *
ATG GCT CGG CCT GGG CAG CGT TGG CTC GGC AAG TGG CTT GTG GCG
Met Ala Arg Pro Gly Gln Arg Trp Leu Gly Lys Trp Leu Val Ala>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

      50      60      70      80      90
      *      *      *      *      *
ATG GTC GTG TGG GCG CTG TGC CGG CTC GCC ACA CCG CTG GCC AAG
Met Val Val Trp Ala Leu Cys Arg Leu Ala Thr Pro Leu Ala Lys>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

     100     110     120     130
      *      *      *      *      *
AAC CTG GAG CCC GTA TCC TGG AGC TCC CTC AAC CCC AAG TTC CTG
Asn Leu Glu Pro Val Ser Trp Ser Ser Leu Asn Pro Lys Phe Leu>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

     140     150     160     170     180
      *      *      *      *      *
AGT GGG AAG GGC TTG GTG ATC TAT CCG AAA ATT GGA GAC AAG CTG
Ser Gly Lys Gly Leu Val Ile Tyr Pro Lys Ile Gly Asp Lys Leu>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

     190     200     210     220
      *      *      *      *      *
GAC ATC ATC TGC CCC CGA GCA GAA GCA GGG CGG CCC TAT GAG TAC
Asp Ile Ile Cys Pro Arg Ala Glu Ala Gly Arg Pro Tyr Glu Tyr>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

     230     240     250     260     270
      *      *      *      *      *
TAC AAG CTG TAC CTG GTG CGG CCT GAG CAG GCA GCT GCC TGT AGC
Tyr Lys Leu Tyr Leu Val Arg Pro Glu Gln Ala Ala Ala Cys Ser>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

     280     290     300     310
      *      *      *      *      *
ACA GTT CTC GAC CCC AAC GTG TTG GTC ACC TGC AAT AGG CCA GAG
Thr Val Leu Asp Pro Asn Val Leu Val Thr Cys Asn Arg Pro Glu>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

     320     330     340     350     360
      *      *      *      *      *
CAG GAA ATA CGC TTT ACC ATC AAG TTC CAG GAG TTC AGC CCC AAC
Gln Glu Ile Arg Phe Thr Ile Lys Phe Gln Glu Phe Ser Pro Asn>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

     370     380     390     400
      *      *      *      *      *
TAC ATG GGC CTG GAG TTC AAG AAG CAC CAT GAT TAC TAC ATT ACC
Tyr Met Gly Leu Glu Phe Lys Lys His His Asp Tyr Tyr Ile Thr>
__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__>

```

31/42

Figure 14B

```

      410      420      430      440      450
      *      *      *      *      *
TCA ACA TCC AAT GGA AGC CTG GAG GGG CTG GAA AAC CGG GAG GGC
Ser Thr Ser Asn Gly Ser Leu Glu Gly Leu Glu Asn Arg Glu Gly>
__a__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__a__>

      460      470      480      490
      *      *      *      *      *
GGT GTG TGC CGC ACA CGC ACC ATG AAG ATC ATC ATG AAG GTT GGG
Gly Val Cys Arg Thr Arg Thr Met Lys Ile Ile Met Lys Val Gly>
__a__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__a__>

      500      510      520      530      540
      *      *      *      *      *
CAA GAT CCC AAT GCT GTG ACG CCT GAG CAG CTG ACT ACC AGC AGG
Gln Asp Pro Asn Ala Val Thr Pro Glu Gln Leu Thr Thr Ser Arg>
__a__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__a__>

      550      560      570      580
      *      *      *      *      *
CCC AGC AAG GAG GCA GAC AAC ACT GTC AAG ATG GCC ACA CAG GCC
Pro Ser Lys Glu Ala Asp Asn Thr Val Lys Met Ala Thr Gln Ala>
__a__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__a__>

      590      600      610      620      630
      *      *      *      *      *
CCT GGT AGT CGG GGC TCC CTG GGT GAC TCT GAT GGC AAG CAT GAG
Pro Gly Ser Arg Gly Ser Leu Gly Asp Ser Asp Gly Lys His Glu>
__a__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__a__>

      640      650      660      670
      *      *      *      *      *
ACT GTG AAC CAG GAA GAG AAG AGT GGC CCA GGT GCA AGT GGG GGC
Thr Val Asn Gln Glu Glu Lys Ser Gly Pro Gly Ala Ser Gly Gly>
__a__a__a__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)__a__a__a__>

      680      690      700      710      720
      *      *      *      *      *
AGC AGC GGG GAC CCT GAT GGC TTC TTC AAC TCC AAG GGC CCG GGT
Ser Ser Gly Asp Pro Asp Gly Phe Phe Asn Ser Lys>
__ELK-L ECTODOMAIN 1 (WITH SIGNAL PEPTIDE)____>
                                     Gly Pro Gly>
                                     __b__b__>

      730      740      750      760
      *      *      *      *      *
AAG AAC CTG GAG CCC GTA TCC TGG AGC TCC CTC AAC CCC AAG TTC
Lys Asn Leu Glu Pro Val Ser Trp Ser Ser Leu Asn Pro Lys Phe>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL)__c__c__c__>

      770      780      790      800      810
      *      *      *      *      *
CTG AGT GGG AAG GGC TTG GTG ATC TAT CCG AAA ATT GGA GAC AAG
Leu Ser Gly Lys Gly Leu Val Ile Tyr Pro Lys Ile Gly Asp Lys>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL)__c__c__c__>

```

32/42
Figure 14C

```

      820      830      840      850
      *      *      *      *
CTG GAC ATC ATC TGC CCC CGA GCA GAA GCA GGG CGG CCC TAT GAG
Leu Asp Ile Ile Cys Pro Arg Ala Glu Ala Gly Arg Pro Tyr Glu>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

      860      870      880      890      900
      *      *      *      *      *
TAC TAC AAG CTG TAC CTG GTG CGG CCT GAG CAG GCA GCT GCC TGT
Tyr Tyr Lys Leu Tyr Leu Val Arg Pro Glu Gln Ala Ala Ala Cys>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

      910      920      930      940
      *      *      *      *      *
AGC ACA GTT CTC GAC CCC AAC GTG TTG GTC ACC TGC AAT AGG CCA
Ser Thr Val Leu Asp Pro Asn Val Leu Val Thr Cys Asn Arg Pro>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

      950      960      970      980      990
      *      *      *      *      *
GAG CAG GAA ATA CGC TTT ACC ATC AAG TTC CAG GAG TTC AGC CCC
Glu Gln Glu Ile Arg Phe Thr Ile Lys Phe Gln Glu Phe Ser Pro>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

     1000     1010     1020     1030
      *      *      *      *      *
AAC TAC ATG GGC CTG GAG TTC AAG AAG CAC CAT GAT TAC TAC ATT
Asn Tyr Met Gly Leu Glu Phe Lys Lys His His Asp Tyr Tyr Ile>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

     1040     1050     1060     1070     1080
      *      *      *      *      *
ACC TCA ACA TCC AAT GGA AGC CTG GAG GGG CTG GAA AAC CGG GAG
Thr Ser Thr Ser Asn Gly Ser Leu Glu Gly Leu Glu Asn Arg Glu>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

     1090     1100     1110     1120
      *      *      *      *      *
GGC GGT GTG TGC CGC ACA CGC ACC ATG AAG ATC ATC ATG AAG GTT
Gly Gly Val Cys Arg Thr Arg Thr Met Lys Ile Ile Met Lys Val>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

     1130     1140     1150     1160     1170
      *      *      *      *      *
GGG CAA GAT CCC AAT GCT GTG ACG CCT GAG CAG CTG ACT ACC AGC
Gly Gln Asp Pro Asn Ala Val Thr Pro Glu Gln Leu Thr Thr Ser>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

     1180     1190     1200     1210
      *      *      *      *      *
AGG CCC AGC AAG GAG GCA GAC AAC ACT GTC AAG ATG GCC ACA CAG
Arg Pro Ser Lys Glu Ala Asp Asn Thr Val Lys Met Ala Thr Gln>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

     1220     1230     1240     1250     1260
      *      *      *      *      *
GCC CCT GGT AGT CGG GGC TCC CTG GGT GAC TCT GAT GCC AAG CAT
Ala Pro Gly Ser Arg Gly Ser Leu Gly Asp Ser Asp Gly Lys His>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

```

33/42
Figure 14D

```

      1270      1280      1290      1300
      *      *      *      *
GAG ACT GTG AAC CAG GAA GAG AAG AGT GGC CCA GGT GCA AGT GGG
Glu Thr Val Asn Gln Glu Glu Lys Ser Gly Pro Gly Ala Ser Gly>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>

      1310      1320      1330      1340      1350
      *      *      *      *      *
GGC AGC AGC GGC GAC CCT GAT GGC TTC TTC AAC TCC AAA GGC CCG
Gly Ser Ser Gly Asp Pro Asp Gly Phe Phe Asn Ser Lys>
__c__c__c__ELK-L ECTODOMAIN 2 (NO SIGNAL) __c__c__c__>
                                           Gly Pro>
                                           __d__>

      1360      1370      1380      1390
      *      *      *      *
GGC GAG CCC AAA TCT TGT GAC AAA ACT CAC ACA TGC CCA CCG TGC
Gly>
__>
      Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys>
      __e__e__e__e__HUMAN IGG1 FC TAG__e__e__e__e__>

      1400      1410      1420      1430      1440
      *      *      *      *      *
CCA GCA CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC CTC TTC CCC
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro>
__e__e__e__e__HUMAN IGG1 FC TAG__e__e__e__e__>

      1450      1460      1470      1480
      *      *      *      *
CCA AAA CCC AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT GAG GTC
Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val>
__e__e__e__e__HUMAN IGG1 FC TAG__e__e__e__e__>

      1490      1500      1510      1520      1530
      *      *      *      *      *
ACA TGC GTG GTG GTG GAC GTG AGC CAC GAA GAC CCT GAG GTC AAG
Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys>
__e__e__e__e__HUMAN IGG1 FC TAG__e__e__e__e__>

      1540      1550      1560      1570
      *      *      *      *
TTC AAC TGG TAC GTG GAC GGC GTG GAG GTG CAT AAT GCC AAG ACA
Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr>
__e__e__e__e__HUMAN IGG1 FC TAG__e__e__e__e__>

      1580      1590      1600      1610      1620
      *      *      *      *      *
AAG CCG CGG GAG GAG CAG TAC AAC AGC ACG TAC CGT GTG GTC AGC
Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser>
__e__e__e__e__HUMAN IGG1 FC TAG__e__e__e__e__>

      1630      1640      1650      1660
      *      *      *      *
GTC CTC ACC GTC CTG CAC CAG GAC TGG CTG AAT GGC AAG GAG TAC
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr>
__c__e__e__e__c__HUMAN IGG1 FC TAG__c__e__e__e__c__>

```

1680 1690 1700 1710
* * * * *
AAG TGC AAG GTC TCC AAC AAA GCC CTC CCA GCC CCC ATC GAG AAA
Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

1720 1730 1740 1750
* * * * *
ACC ATC TCC AAA GCC AAA GGG CAG CCC CGA GAA CCA CAG GTG TAC
Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

1760 1770 1780 1790 1800
* * * * *
ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC AAG AAC CAG GTC AGC
Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

1810 1820 1830 1840
* * * * *
CTG ACC TGC CTG GTC AAA GGC TTC TAT CCC AGC GAC ATC GCC GTG
Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

1850 1860 1870 1880 1890
* * * * *
GAG TGG GAG AGC AAT GGG CAG CCG GAG AAC AAC TAC AAG ACC ACG
Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

1900 1910 1920 1930
* * * * *
CCT CCC GTG CTG GAC TCC GAC GGC TCC TTC TTC CTC TAC AGC AAG
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

1940 1950 1960 1970 1980
* * * * *
CTC ACC GTG GAC AAG AGC AGG TGG CAG CAG GGG AAC GTC TTC TCA
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

1990 2000 2010 2020
* * * * *
TGC TCC GTG ATG CAT GAG GCT CTG CAC AAC CAC TAC ACG CAG AAG
Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

2030 2040 2050
* * * * *
AGC CTC TCC CTG TCT CCG GGT AAA TGA
Ser Leu Ser Leu Ser Pro Gly Lys ***>
_e_e_e_e_e_HUMAN IGG1 FC TAG_e_e_e_e_e_e_e_>

35/42
Figure 15A

```

      10      20      30      40
      *      *      *      *
ATG GCC ATG GCC CGG TCC AGG AGG GAC TCT GTG TGG AAG TAC TGT
Met Ala Met Ala Arg Ser Arg Arg Asp Ser Val Trp Lys Tyr Cys>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

      50      60      70      80      90
      *      *      *      *      *
TGG GGA CTT TTG ATG GTT TTG TGC AGA ACT GCG ATC TCC AGA TCG
Trp Gly Leu Leu Met Val Leu Cys Arg Thr Ala Ile Ser Arg Ser>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

     100     110     120     130
     *      *      *      *      *
ATA GTT TTA GAG CCT ATC TAC TGG AAT TCC TCG AAC TCC AAA TTT
Ile Val Leu Glu Pro Ile Tyr Trp Asn Ser Ser Asn Ser Lys Phe>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

     140     150     160     170     180
     *      *      *      *      *
CTA CCC GGA CAA GGC CTG GTA CTA TAC CCA CAG ATA GGA GAC AAA
Leu Pro Gly Gln Gly Leu Val Leu Tyr Pro Gln Ile Gly Asp Lys>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

     190     200     210     220
     *      *      *      *      *
TTG GAT ATT ATT TGC CCC AAA GTG GAC TCT AAA ACT GTT GGC CAG
Leu Asp Ile Ile Cys Pro Lys Val Asp Ser Lys Thr Val Gly Gln>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

     230     240     250     260     270
     *      *      *      *      *
TAT GAA TAT TAT AAA GTT TAT ATG GTT GAT AAA GAC CAA GCA GAC
Tyr Glu Tyr Tyr Lys Val Tyr Met Val Asp Lys Asp Gln Ala Asp>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

     280     290     300     310
     *      *      *      *      *
AGA TGC ACA ATT AAG AAG GAG AAT ACC CCG CTG CTC AAC TGT GCC
Arg Cys Thr Ile Lys Lys Glu Asn Thr Pro Leu Leu Asn Cys Ala>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

     320     330     340     350     360
     *      *      *      *      *
AGA CCA GAC CAA GAT GTG AAA TTC ACC ATC AAG TTT CAA GAA TTC
Arg Pro Asp Gln Asp Val Lys Phe Thr Ile Lys Phe Gln Glu Phe>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

     370     380     390     400
     *      *      *      *      *
AGC CCT AAC CTC TGG GGT CTA GAA TTT CAG AAG AAC AAA GAT TAC
Ser Pro Asn Leu Trp Gly Leu Glu Phe Gln Lys Asn Lys Asp Tyr>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

```

36/42
Figure 15B

```

      410      420      430      440      450
      *      *      *      *      *
TAC ATT ATA TCT ACA TCA AAT GGG TCT TTG GAG GGC CTG GAT AAC
Tyr Ile Ile Ser Thr Ser Asn Gly Ser Leu Glu Gly Leu Asp Asn>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

      460      470      480      490
      *      *      *      *      *
CAG GAG GGA GGG CTG TCC CAG ACA AGA GCC ATG AAG ATC CTC ATG
Gln Glu Gly Gly Val Cys Gln Thr Arg Ala Met Lys Ile Leu Met>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

      500      510      520      530      540
      *      *      *      *      *
AAA GTT GGA CAA GAT GCA AGT TCT GCT GGA TCA GCC AGG AAT CAC
Lys Val Gly Gln Asp Ala Ser Ser Ala Gly Ser Ala Arg Asn His>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

      550      560      570      580
      *      *      *      *      *
GGT CCA ACA AGA CGT CCA GAG CTA GAA GCT GGT ACA AAT GGG AGA
Gly Pro Thr Arg Arg Pro Glu Leu Glu Ala Gly Thr Asn Gly Arg>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

      590      600      610      620      630
      *      *      *      *      *
AGT TCA ACA ACA AGT CCC TTT GTG AAG CCA AAT CCA GGT TCT AGC
Ser Ser Thr Thr Ser Pro Phe Val Lys Pro Asn Pro Gly Ser Ser>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

      640      650      660      670
      *      *      *      *      *
ACC GAT GGC AAC AGC GCG GGG CAT TCC GGG AAC AAT CTC CTG GGG
Thr Asp Gly Asn Ser Ala Gly His Ser Gly Asn Asn Leu Leu Gly>
__a__EPHRIN-B2  ECTO DOMAIN 1 (WITH SIGNAL PEPTIDE) __a__>

      680      690      700      710      720
      *      *      *      *      *
GGC CCG GGA ATA GTT TTA GAG CCT ATC TAC TGG AAT TCC TCG AAC
Gly Pro Gly>
__b__b__>
      Ile Val Leu Glu Pro Ile Tyr Trp Asn Ser Ser Asn>
      __EPHRIN-B2  ECTO DOMAIN 2 ( WITHOUT SIGNA__>

      730      740      750      760
      *      *      *      *      *
TCC AAA TTT CTA CCC GGA CAA GGC CTG GTA CTA TAC CCA CAG ATA
Ser Lys Phe Leu Pro Gly Gln Gly Leu Val Leu Tyr Pro Gln Ile>
__EPHRIN-B2  ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE) __e__>

      770      780      790      800      810
      *      *      *      *      *
GGA GAC AAA TTG GAT ATT ATT TGC CCC AAA GTG GAC TCT AAA ACT
Gly Asp Lys Leu Asp Ile Ile Cys Pro Lys Val Asp Ser Lys Thr>
__EPHRIN-B2  ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE) __c__>

```

37/42
Figure 15C

```

      820      830      840      850
      *      *      *      *
GTT GGC CAG TAT GAA TAT TAT AAA GTT TAT ATG GTT GAT AAA GAC
Val Gly Gln Tyr Glu Tyr Tyr Lys Val Tyr Met Val Asp Lys Asp>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      860      870      880      890      900
      *      *      *      *      *
CAA GCA GAC AGA TGC ACA ATT AAG AAG GAG AAT ACC CCG CTG CTC
Gln Ala Asp Arg Cys Thr Ile Lys Lys Glu Asn Thr Pro Leu Leu>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      910      920      930      940
      *      *      *      *
AAC TGT GCC AGA CCA GAC CAA GAT GTG AAA TTC ACC ATC AAG TTT
Asn Cys Ala Arg Pro Asp Gln Asp Val Lys Phe Thr Ile Lys Phe>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      950      960      970      980      990
      *      *      *      *      *
CAA GAA TTC AGC CCT AAC CTC TGG GGT CTA GAA TTT CAG AAG AAC
Gln Glu Phe Ser Pro Asn Leu Trp Gly Leu Glu Phe Gln Lys Asn>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      1000     1010     1020     1030
      *      *      *      *
AAA GAT TAC TAC ATT ATA TCT ACA TCA AAT GGG TCT TTG GAG GGC
Lys Asp Tyr Tyr Ile Ile Ser Thr Ser Asn Gly Ser Leu Glu Gly>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      1040     1050     1060     1070     1080
      *      *      *      *      *
CTG GAT AAC CAG GAG GGA GGG GTG TGC CAG ACA AGA GCC ATG AAG
Leu Asp Asn Gln Glu Gly Gly Val Cys Gln Thr Arg Ala Met Lys>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      1090     1100     1110     1120
      *      *      *      *
ATC CTC ATG AAA GTT GGA CAA GAT GCA AGT TCT GCT GGA TCA GCC
Ile Leu Met Lys Val Gly Gln Asp Ala Ser Ser Ala Gly Ser Ala>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      1130     1140     1150     1160     1170
      *      *      *      *      *
AGG AAT CAC GGT CCA ACA AGA CGC CCA GAG CTA GAA GCT GGT ACA
Arg Asn His Gly Pro Thr Arg Arg Pro Glu Leu Glu Ala Gly Thr>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      1180     1190     1200     1210
      *      *      *      *
AAT GGG AGA AGT TCA ACA ACA AGT CCC TTT GTG AAG CCA AAT CCA
Asn Gly Arg Ser Ser Thr Thr Ser Pro Phe Val Lys Pro Asn Pro>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

      1220     1230     1240     1250     1260
      *      *      *      *      *
GGT TCT AGC ACC GAT GGC AAC AGC GCG GGC CAT TCC GGG AAC AAT
Gly Ser Ser Thr Asp Gly Asn Ser Ala Gly His Ser Gly Asn Asn>
____EPHRIN-B2 ECTO DOMAIN 2 ( WITHOUT SIGNAL PEPTIDE)_e____>

```

38/42
Figure 15D

```

      1270      1280      1290      1300
      *      *      *      *      *
CTC CTG GGG G GC CCG GGC GAG CCC AAA TCT TGT GAC AAA ACT CAC
      Glu Pro Lys Ser Cys Asp Lys Thr His>
      ____c____HUMAN IGG1 FC TAG_c__c__>
      Gly Pro Gly>
      _d_d_d_d_>
Leu Leu Gly Xxx>
__e__e__e__>

      1310      1320      1330      1340      1350
      *      *      *      *      *
ACA TGC CCA CCG TGC CCA GCA CCT GAA CTC CTG GGG GGA CCG TCA
Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

      1360      1370      1380      1390
      *      *      *      *      *
GTC TTC CTC TTC CCC CCA AAA CCC AAG GAC ACC CTC ATG ATC TCC
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

      1400      1410      1420      1430      1440
      *      *      *      *      *
CGG ACC CCT GAG GTC ACA TGC GTG GTG GTG GAC GTG AGC CAC GAA
Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

      1450      1460      1470      1480
      *      *      *      *      *
GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG GAG GTG
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

      1490      1500      1510      1520      1530
      *      *      *      *      *
CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG TAC AAC AGC ACG
His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

      1540      1550      1560      1570
      *      *      *      *      *
TAC CGT GTG GTC AGC GTC CTC ACC GTC CTG CAC CAG GAC TGG CTG
Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

      1580      1590      1600      1610      1620
      *      *      *      *      *
AAT GGC AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA GCC CTC CCA
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

      1630      1640      1650      1660
      *      *      *      *      *
CCC CCC ATC GAG AAA ACC ATC TCC AAA GCC AAA GGG CAG CCC CGA
Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Glu Pro Arg>
__c__c__c__c__c__HUMAN IGG1 FC TAG_c__c__c__c__c__>

```

1670 1680 1690 1700 1710
GAA CCA CAG GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC
Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr>
_c_c_c_c_c_____HUMAN IGG1 FC TAG_c_c_c_c_c____>

1720 1730 1740 1750
AAG AAC CAG GTC AGC CTG ACC TGC CTG GTC AAA GGC TTC TAT CCC
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro>
_c_c_c_c_c_____HUMAN IGG1 FC TAG_c_c_c_c_c____>

1760 1770 1780 1790 1800
AGC GAC ATC GCC GTG GAG TGG GAG AGC AAT GGG CAG CCG GAG AAC
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn>
_c_c_c_c_c_____HUMAN IGG1 FC TAG_c_c_c_c_c____>

1810 1820 1830 1840
AAC TAC AAG ACC ACG CCT CCC GTG CTG GAC TCC GAC GGC TCC TTC
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe>
_c_c_c_c_c_____HUMAN IGG1 FC TAG_c_c_c_c_c____>


1850 1860 1870 1880 1890
TTC CTC TAC AGC AAG CTC ACC GTG GAC AAG AGC AGG TGG CAG CAG
Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln>
_c_c_c_c_c_____HUMAN IGG1 FC TAG_c_c_c_c_c____>

1900 1910 1920 1930
GGG AAC GTC TTC TCA TGC TCC GTG ATG CAT GAG GCT CTG CAC AAC
Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn>
_c_c_c_c_c_____HUMAN IGG1 FC TAG_c_c_c_c_c____>

1940 1950 1960 1970
CAC TAC ACG CAG AAG AGC CTC TCC CTG TCT CCG GGT AAA TGA
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys ***>
_c_c_c_c_c_____HUMAN IGG1 FC TAG_c_c_c_c_c____>


40/42

Figure 16




41/42

Figure 17



42/42

Figure 18



1 2 3 4 5 6 7 8 9

INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/US 99/30900

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N15/62 C12N5/10 C12N1/21 C07K14/515
C07K14/52

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 37621 A (MORPHOSYS PROTEINOPTIMIERUNG ;PACK PETER (DE); HOESS ADOLF (DE)) 28 November 1996 (1996-11-28) abstract page 1, line 12 - line 15 page 2, line 4 - line 9 page 14, line 6 - line 11 page 16, line 29 - line 34 figure 1A	1-5, 12-22
Y A	---	8-11 6,7
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 May 2000

Date of mailing of the international search report

09/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 99/30900

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 03569 A (SANGSTAT MEDICAL CORP) 5 March 1992 (1992-03-05) abstract page 1 -page 3 page 19, line 14 - line 23	23-26, 29,32-41
Y		
A		30 27,28
Y	WO 95 27060 A (REGENERON PHARMA) 12 October 1995 (1995-10-12) cited in the application abstract page 23, line 21 -page 24, line 8 claims 4,8	8-11,30
A	EP 0 816 510 A (TORAY RESEARCH CENTER INC ;TORAY INDUSTRIES (JP)) 7 January 1998 (1998-01-07) abstract	1-22
A	DAVIS S. ET AL.: "Isolation of Angiopoietin-1, a ligand for the TIE2 receptor, by secretion-trap expression cloning" CELL, vol. 87, 27 December 1996 (1996-12-27), pages 1161-1169, XP002138354 cited in the application the whole document	6,7,27, 28
A	MAISONPIERRE P.C. ET AL.: "Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis." SCIENCE, vol. 277, 4 July 1997 (1997-07-04), pages 55-60, XP002138355 cited in the application the whole document	6,7,27, 28
A	TOURNAY C. ET AL.: "Uptake of recombinant myeloperoxidase, free or fused to Fc-gamma, by macrophages enhances killing activity towards micro-organisms" DNA CELL BIOLOGY, vol. 15, no. 8, 1996, pages 617-624, XP000907279 abstract page 618	21,40

INTERNATIONAL SEARCH REPORT

Information on patent family members

In ternational Application No
PCT/US 99/30900

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9637621 A	28-11-1996	CA 2222055 A EP 0827544 A JP 11508126 T	28-11-1996 11-03-1998 21-07-1999
W0 9203569 A	05-03-1992	CA 2090105 A EP 0547163 A JP 6502301 T US 5672486 A	01-03-1992 23-06-1993 17-03-1994 30-09-1997
W0 9527060 A	12-10-1995	US 5747033 A AU 691915 B AU 2278995 A CA 2187167 A EP 0758381 A JP 9511401 T ZA 9502762 A	05-05-1998 28-05-1998 23-10-1995 12-10-1995 19-02-1997 18-11-1997 20-02-1996
EP 0816510 A	07-01-1998	CA 2213512 A W0 9723639 A	03-07-1997 03-07-1997

THIS PAGE BLANK (USPTO)